NT 00 11		
Name of Subject: Gender:		
Dear Sir/Madam:		
The result of the latent tu interferon-gamma release a you have LTBI, but not act can give more than 90%	berculosis infection (LTBI) test (Tuberculin ssay, IGRA), and the evaluation by the phy ive tuberculosis (TB). A complete course of protection from developing active TB dissibility of onset of TB disease and further	sician indicated that treatment for LTBI ease. This helps to
Special Notice of LTBI tro	eatment:	
I. The following regimens a	are recommended for the treatment of LTBI.	(Please mark V)
	laily isoniazid plus rifapentine (28 doses)	
☐3HP: Three months of	of once-weekly isoniazid plus rifapentine (12	2 doses)
\Box 4R: Four months of c	laily rifampin (120 doses)	
☐3HR: Three months of	of daily isoniazid plus rifampin (90 doses)	
☐6H: Six months of da	tily isoniazid (180 doses)	
☐9H: Nine months of o	daily isoniazid (270 doses)	
others:		
•	eventive therapy (DOPT) observer will be u complete the treatment.	arranged to provide
treatment. During the tre loss of appetite, pain in te eyes, rashes, acute alle	nical evaluation at least once a month duatment, if you do not feel well (such as feve the upper right abdomen, numbness of limb rgy, and other symptoms). Please informmanagers or the physicians in hospitals at or	r, dizziness, nausea, s, yellowish skin or n the public health
IV. If you have any questio number	ns, please do not hesitate to contact us at pho-	one
Public Health Bureau to s	public health station under thesafeguard my medications on LTBI, and health workers for DOPT service.	
LTB	Treatment and DOPT Consent	
With regard to the declaration above:	Relation between the undersign subject of treatment	ned and the
☐ Agree	☐ Myself	_(signature)
☐ Disagree	☐ Appointed Guardian	(signature)