



## Summary : Week 22 (May 26–June 1, 2013)

During week 22, 2013, influenza activity in Taiwan is still at low to moderate level. From the virologic surveillance data, the proportion of influenza A (H1N1) viruses has been co-circulating with influenza A (H3N2) viruses in the community since week 8.

- Of the 149 specimens tested during week 20, 2013, 29 (19.5%) were positive for influenza viruses, 18 (12.1%) were positive for influenza A (H1N1) virus, 10 (6.7%) were positive for influenza A (H3N2) virus, and 1 (0.7%) were positive for influenza B virus.
- During week 22, 2013, there were 43 new cases of complicated influenza, including 28 cases with influenza A (H3N2) virus infection, 14 cases with influenza A (H1N1) virus infection; no report of death from complicated influenza. Since July 1, 2012, 825 cases of complicated influenza infection have been confirmed, including 525 cases from influenza A (H3N2) virus infection, 283 cases from influenza A (H1N1) virus infection, 11 cases from influenza B virus infection, and 6 cases due to un-typed influenza A virus infection. 245 cases of complicated influenza infection received the intensive care. Since July 1, 2012, there have been 60 reports of death from complicated influenza infection; 36 cases from influenza A (H3N2) virus infection and 24 cases from influenza A (H1N1) virus infection, respectively.
- During week 21, 2013, the number of deaths related to pneumonia and influenza was 257, which was above the number of the previous week (253 deaths).
- The proportion of outpatient visits for influenza-like illness (ILI) is 1.09%, a 4.4% decrease was compared with the proportion of the previous week (1.13%).
- The proportion of emergency room visits for ILI is 10.59%, which was comparable with the proportion of the previous week (10.59%).

## Virologic surveillance


During week 20, 2013, the results of tests performed are summarized in the table below.

	Data for week 20	Cumulative data since 7/1/2012
Number of specimens tested	149	7,385
Number of positive specimens (%)	29(19.5)	921(12.5)
Positive specimens by type/subtype (%)		
Influenza A (% of all positive specimens)	28(97)	904(98)
A (H1N1) (% of all Influenza A)	18(64)	300(33)
A (H3N2)	10(36)	602(67)
A (unable to subtype)	0(0)	2(<1)
A (subtyping not performed)	0(0)	0(0)
Influenza B	1(3)	17(2)



**Antigenic characterization:** Taiwan CDC has antigenically characterized 207 human influenza viruses [85 influenza A (H1N1) viruses, 109 influenza A (H3N2) viruses, and 13 influenza B viruses] since October 1, 2012. 96% (n=105) of the influenza A (H3N2) viruses tested were related to the A (H3N2) component of the 2012-13 influenza vaccine (A/Victoria/361/2011(H3N2)). 99% (n=84) influenza A (H1N1) viruses tested were related to the A (H1N1) component of the 2012-13 influenza vaccine (A/California/7/2009(H1N1)pdm09). 77% (n=10) of the influenza B viruses tested belonged to the B/Yamagata lineage and were related to the B component of the 2012-13 influenza vaccine (B/Wisconsin/1/2010).

### Influenza positive tests reported to Taiwan CDC by contracted laboratories, 2011–2013



**Antiviral resistance:** Since October 1, 2012, 42 influenza A (H1N1), 66 influenza A (H3N2) and 2 influenza B viruses have been tested for resistance to the neuraminidase inhibitors (oseltamivir). The results of antiviral resistance testing performed on these viruses are summarized in the table below.


	Isolates tested (n)	Resistance Viruses, n (%)
		Oseltamivir
Influenza A (H1N1)	42	0 (0)
Influenza A (H3N2)	66	0 (0)
Influenza B	2	0 (0)






## Pneumonia and influenza (P&I) mortality surveillance

During week 21, 2013, 257 deaths reported through the National Death Certificate System mentioned P&I as the cause of death. This number was above the number of the previous week (253 deaths). The number of deaths related to P&I for adults of 65 or greater is the highest among the three age groups (0–49, 50–64, and 65+).

### National pneumonia and influenza mortality Week ending May 25, 2013




## National pneumonia and influenza mortality by age group Week ending May 25, 2013



## Reports of complicated influenza\*


During week 22, 2013, there were 43 new cases of complicated influenza, including 28 cases with influenza A (H3N2) virus infection, 14 cases with influenza A (H1N1) virus infection, and 1 case with influenza B virus infection; no report of death from complicated influenza. Since July 1, 2012, 825 cases of complicated influenza infection have been confirmed, including 525 cases from influenza A (H3N2) virus infection, 283 cases from influenza A (H1N1) virus infection, 11 cases from influenza B virus infection, and 6 cases due to un-typed influenza A virus infection. 245 cases of complicated influenza infection received the intensive care. Since July 1, 2012, there have been 60 reports of death from complicated influenza infection; 36 cases from influenza A (H3N2) virus infection and 24 cases from influenza A (H1N1) virus infection, respectively.

### Number of complicated influenza reports by week of onset June 26, 2011 to present



\*Defined as influenza infection with pulmonary complication, neurologic complication, myocarditis, pericarditis, invasive bacterial infection requiring hospitalization, intensive care or resulting in deaths.

### Rate of complicated influenza reports by age groups July 1, 2012 to present



\*Numbers represent number of complicated influenza reports for that specific age stratum.



## Outpatient and emergency room influenza-like illness surveillance

Nationwide during week 22, 2013, the proportion of outpatient visits for ILI is 1.09%, a 4.4% decrease was compared with the proportion of the previous week (1.13%).

The proportion of emergency room visits for ILI is 10.59%, which was comparable with the proportion of the previous week (10.59%).

**Proportions of outpatient and emergency room (ER) visits for influenza-like illness (June 26, 2011 to present)**

