



Summary : Week 8 (Feb. 19– Feb. 25, 2012)

Influenza activity in Taiwan peaked at week 2. Seasonal influenza virus type B is predominant in the community.

- Of the 217 specimens tested during week 6, 2012, 72 (33.2%) were positive for influenza viruses, 59 (27.2%) were positive for influenza B viruses, 7 (3.2%) were positive for influenza AH3 viruses, 6 (2.8%) were positive for pandemic (H1N1) 2009 virus.
- There were 50 new complicated influenza infections and no report of death from complicated influenza infection during week 8. Since July 1, 2011, 1173 complicated influenza infections have been confirmed, including 930 cases from influenza B virus infection, 185 cases from influenza A virus infection, 45 cases from pandemic (H1N1) 2009 virus infection and 13 cases due to un-typed influenza A virus infection. 395 cases of complicated influenza infections received the intensive care. Since July 1, 2011, there have been 72 reports of death from complicated influenza infection; 57 from the seasonal influenza B strain, 13 from the seasonal H3N2 strain, 1 from pandemic (H1N1) 2009 virus infection and 1 from the un-typed influenza A virus, respectively.
- The number of deaths related to pneumonia and influenza during week 7 was 361, which was below the number of the previous week (384 deaths).
- The rate of outpatient visits for influenza-like illness is 1.68%, a 2.3% decrease compared with the rate of previous week (1.72%).
- The rate of emergency room visits for ILI is 12.97%, a 4.8 % decrease compared with the rate of previous week (13.62%).

Virologic surveillance


The results of tests performed during week 6 are summarized in the table below.

	Data for week 6	Cumulative data since 7/1/2011
Number of specimens tested	217	12218
Number of positive specimens (%)	72(33)	2661(22)
Positive specimens by type/subtype (%)		
Influenza A (% of all positive specimens)	13(18)	292(11)
A (pandemic [H1N1] 2009) (% of all Influenza A)	6(46)	49(17)
A (H3)	7(54)	242(83)
A (H1)	0(0)	0(0)
A (unable to subtype)	0(0)	1(0.3)
A (subtyping not performed)	0(0)	0(0)
Influenza B	59(82)	2369(89)



Antigenic characterization: Taiwan CDC has antigenically characterized 326 human influenza viruses [2 pandemic (H1N1) 2009, 36 influenza A (H3), 288 influenza B] since July 1, 2011. 94.4% (n=34) of the influenza A (H3N2) viruses tested were related to the A (H3N2) vaccine component (A/Perth/16/2009). 13.5% (n=39) of the influenza B viruses tested belonged to the B/Victoria lineage and were related to the B component of the 2011-12 influenza vaccine (B/Brisbane/60/2008).

Influenza positive tests reported to Taiwan CDC by contracted laboratories, 2010–2012



Antiviral resistance: Since July 1, 2011, 7 pandemic (H1N1) 2009, 70 influenza A (H3N2) and 198 influenza B viruses have been tested for resistance to the neuraminidase inhibitors (oseltamivir). The results of antiviral resistance testing performed on these viruses are summarized in the table below.


	Isolates tested (n)	Resistance Viruses, n (%)
		Oseltamivir
A (pandemic [H1N1] 2009)	7	0(0)
A (H3)	70	0 (0)
B	198	0 (0)





Pneumonia and influenza (P&I) mortality surveillance

During week 7, 2012, 361 deaths reported through the National Death Certificate System mentioned P&I as the cause of death. This number was below the number of previous week (384 deaths). The number of deaths related to P&I for age groups 0–49, 50–64, and greater than 65 years was the highest for adult greater than 65 years of age.

National pneumonia and influenza mortality Week ending 18/02/2012




National pneumonia and influenza mortality by age group Week ending 18/02/2012



Reports of complicated influenza*


There were 50 new complicated influenza infections (including 39 seasonal influenza B cases, 6 seasonal influenza AH3 cases, 3 cases from pandemic (H1N1) 2009 virus infection and 2 cases due to un-typed influenza A virus infection) during week 8. Since July 1, 2011, 1173 complicated influenza infections have been confirmed, including 395 cases requiring intensive care. Since July 1, 2011, there have been 72 reports of death from complicated influenza infection; 57 from the seasonal influenza B strain, 13 from the seasonal H3N2 strain, 1 from pandemic (H1N1) 2009 virus and 1 from the un-typed influenza A virus, respectively.

Number of complicated influenza reports by week of onset 27/6/2010 to present



*Defined as influenza infection with pulmonary complication, neurologic complication, myocarditis, pericarditis, invasive bacterial infection requiring hospitalization, intensive care or resulting in deaths.

Rate of complicated influenza reports by age groups 1/7/2011 to present



*Numbers represent number of complicated influenza reports for that specific age stratum.



Outpatient and emergency room influenza-like illness surveillance

Nationwide during week 8, 1.68% of outpatient visits reported through the National Health Insurance Database were due to influenza-like illness (ILI), a 2.3% decrease compared with the rate of previous week (1.72%).

The rate of emergency room visits for ILI is 12.97%, a 4.8% decrease compared with the rate of previous week (13.62%).

Rate of outpatient and emergency room (ER) visits for influenza-like illness

27/6/2010 to present

