



Summary : Week 15 (Apr. 7–Apr. 13, 2013)

During week 15, 2013, influenza activity in Taiwan is still at low to moderate level. From the virologic surveillance data, the proportion of influenza A (H1N1) viruses has been increasing and co-circulating with influenza A (H3N2) viruses in the community since week 8.

- Of the 221 specimens tested during week 13, 2013, 45 (20.4%) were positive for influenza viruses, 24 (10.9%) were positive for influenza A (H1N1) virus, 20 (9%) were positive for influenza A (H3N2) virus, and 1 (0.5%) was positive for influenza B virus.
- During week 15, 2013, there were 43 new cases of complicated influenza, including 17 cases with influenza A (H3N2) virus infection, 25 cases with influenza A (H1N1) virus infection, and 1 case with un-typed influenza A virus infection; 2 reports of death from complicated influenza with influenza A (H1N1) virus infection. Since July 1, 2012, 574 cases of complicated influenza infection have been confirmed, including 393 cases from influenza A (H3N2) virus infection, 168 cases from influenza A (H1N1) virus infection, 7 cases due to un-typed influenza A virus infection, and 6 cases from influenza B virus infection. 176 cases of complicated influenza infection received the intensive care. Since July 1, 2012, there have been 43 reports of death from complicated influenza infection; 30 cases from influenza A (H3N2) virus infection and 13 cases from influenza A (H1N1) virus infection, respectively.
- During week 14, 2013, the number of deaths related to pneumonia and influenza was 259, which was below the number of the previous week (269 deaths).
- The proportion of outpatient visits for influenza-like illness is 1.32%, a 3.6% decrease compared with the proportion of previous week (1.37%).
- The proportion of emergency room visits for ILI is 11.71%, a 4.1% decrease compared with the proportion of previous week (12.21%).

Virologic surveillance


During week 13, 2013, the results of tests performed are summarized in the table below.

	Data for week 13	Cumulative data since 7/1/2012
Number of specimens tested	221	6044
Number of positive specimens (%)	45(20.4)	661(10.9)
Positive specimens by type/subtype (%)		
Influenza A (% of all positive specimens)	44(98)	648(98)
A (H1N1) (% of all Influenza A)	24(55)	149(23)
A (H3N2)	20(45)	497(77)
A (unable to subtype)	0(0)	2(<1)
A (subtyping not performed)	0(0)	0(0)
Influenza B	1(2)	13(2)



Antigenic characterization: Taiwan CDC has antigenically characterized 83 human influenza viruses [25 influenza A (H1N1) viruses, 52 influenza A (H3N2) viruses, and 6 influenza B viruses] since October 1, 2012. 98% (n=51) of the influenza A (H3N2) viruses tested were related to the A (H3N2) component of the 2012-13 influenza vaccine (A/Victoria/361/2011(H3N2)). All of the 25 influenza A (H1N1) viruses tested were related to the A (H1N1) component of the 2012-13 influenza vaccine (A/California/7/2009(H1N1)pdm09). 83% (n=5) of the influenza B viruses tested belonged to the B/Yamagata lineage and were related to the B component of the 2012-13 influenza vaccine (B/Wisconsin/1/2010).

Influenza positive tests reported to Taiwan CDC by contracted laboratories, 2011–2013



Antiviral resistance: Since October 1, 2012, 34 influenza A (H1N1), 60 influenza A (H3N2) and 2 influenza B viruses have been tested for resistance to the neuraminidase inhibitors (oseltamivir). The results of antiviral resistance testing performed on these viruses are summarized in the table below.


	Isolates tested (n)	Resistance Viruses, n (%)
		Oseltamivir
Influenza A (H1N1)	34	0 (0)
Influenza A (H3N2)	60	0 (0)
Influenza B	2	0 (0)






Pneumonia and influenza (P&I) mortality surveillance

During week 14, 2013, 259 deaths reported through the National Death Certificate System mentioned P&I as the cause of death. This number was below the number of previous week (269 deaths). The number of deaths related to P&I for adults of 65 or greater is the highest among the three age groups (0–49, 50–64, and 65⁺).

National pneumonia and influenza mortality Week ending April 6, 2013




National pneumonia and influenza mortality by age group Week ending April 6, 2013



Reports of complicated influenza*


During week 15, 2013, there were 43 new cases of complicated influenza, including 17 cases with influenza A (H3N2) virus infection, 25 cases with influenza A (H1N1) virus infection, and 1 case with un-typed influenza A virus infection; 2 reports of death from complicated influenza with influenza A (H1N1) virus infection. Since July 1, 2012, 574 cases of complicated influenza infection have been confirmed, including 393 cases from influenza A (H3N2) virus infection, 168 cases from influenza A (H1N1) virus infection, 7 cases due to un-typed influenza A virus infection, and 6 cases from influenza B virus infection. 176 cases of complicated influenza infection received the intensive care. Since July 1, 2012, there have been 43 reports of death from complicated influenza infection; 30 cases from influenza A (H3N2) virus infection and 13 cases from influenza A (H1N1) virus infection, respectively.

Number of complicated influenza reports by week of onset June 26, 2011 to present



*Defined as influenza infection with pulmonary complication, neurologic complication, myocarditis, pericarditis, invasive bacterial infection requiring hospitalization, intensive care or resulting in deaths.

Rate of complicated influenza reports by age groups July 1, 2012 to present



*Numbers represent number of complicated influenza reports for that specific age stratum.



Outpatient and emergency room influenza-like illness surveillance

Nationwide during week 15, 2013, 1.32% of outpatient visits reported through the National Health Insurance Database were due to influenza-like illness (ILI), a 3.6% decrease compared with the proportion of previous week (1.37%).

The proportion of emergency room visits for ILI is 11.71%, a 4.1% decrease compared with the proportion of previous week (12.21%).

Proportions of outpatient and emergency room (ER) visits for influenza-like illness (June 26, 2011 to present)

