



Synopsis

Influenza activity decreased but remained elevated during week 5. The most frequently identified influenza virus type was influenza B.


- The proportions of ILI visit in outpatient department and ER decreased compared to the previous week.
- The number of influenza positive specimens was lower than the previous week. The predominant isolated influenza virus was influenza B/Yamagata.
- The number of newly reported influenza cases with severe complications was slightly lower, but the newly confirmed cases remained similar compared to the previous week. There were 53 newly confirmed severe complicated influenza cases and 9 newly fatal cases. A total of 314 severe complicated influenza cases were confirmed since October 1, 2017, and 49 of them were fatal. Influenza B was the predominant virus strain among severe cases and fatal cases.

Viral Surveillance

Types and Trend

According to LARS¹, the number of influenza positive specimens during week 5 was lower than the previous week, and the majority of positive specimens was influenza B virus. The proportion of positive specimens for influenza B virus was 75% during week 5.

Trend of influenza positive specimens according to LARS




¹ In order to present the trend of influenza virus in real-time, the Automated Laboratory Reporting System (LARS) has been established by Taiwan CDC since 2014. The data presented here collected from 51 participating hospitals. All positive specimens data uploads to LARS automatically.



According to the Taiwan CDC Contracted Virology Laboratories², the proportion of influenza positive specimens was 44.7%. Among these, 91.8% were influenza B virus during week 3, 2018. Weekly virus data are available at: <http://nidss.cdc.gov.tw/>.

Influenza isolates and positive rate according to Contracted Virology Laboratories July 1, 2016 to January 20, 2018



Antigenicity

In the past 4 weeks, among the influenza isolates were antigenically characterized, all (100%) of the influenza A (H1N1) virus isolates matched the A (H1N1) component of the 2017-18 influenza vaccine (A/Michigan/45/2015), and 80% of the H3N2 virus isolates matched the A (H3N2) component of the 2017-18 influenza vaccine (A/Hong Kong/4801/2014). Among influenza B isolates, 2% were B/Victoria lineage, and 100% of those isolates matched the B component of the 2017-18 influenza vaccine B/Brisbane/60/2008; 98% were B/Yamagata lineage, and 100% of those isolates matched the B component of the 2017-18 influenza vaccine B/Phuket/3073/2013 (tetraivalent).

Antiviral Resistance

The table below summarized antiviral resistance to neuraminidase inhibitor (Oseltamivir) from October 1, 2017 to present. All of the influenza isolates were susceptible to Oseltamivir.


	Isolates tested (n)	Resistance Viruses, n (%)
		Oseltamivir
Influenza A (H1N1)	14	0
Influenza A (H3N2)	49	0
Influenza B	111	0

² The Contracted Virology Laboratories, including 8 laboratories of medical centers, have been established by Taiwan CDC since March, 1999 to monitor the subtype, antigenicity and drug resistance of influenza viruses in the community.



Influenza-like Illness (ILI) Surveillance

During week 5, the proportion of the outpatient department visits for ILI was 1.97%, and the proportion of ER visits for ILI was 13.85%. Both proportions were lower than the previous week.



* Since 2016, the analysis of the ILI data from National Health Insurance Database is based on the ICD-10 diagnosis codes.


Severe Complicated Influenza Report

The number of newly reported influenza cases with severe complications was slightly lower, but the newly confirmed cases remained similar compared to the previous week. There were 53 newly confirmed cases (45 were influenza B, 5 were H1N1, 3 were H3N2) and 9 newly fatal cases with influenza B virus infection.

In the previous influenza outbreak, the activity returned to the baseline in mid-August 2017 and the number of severe cases continuously declined until September. Since October 1, 2017, a total of 314 severe complicated influenza cases were confirmed, and 49 of them were fatal [the majority of detected virus was influenza B (about 80%), followed by influenza A/H3N2 (about 10%)]. Among these cases, incidence and mortality were the highest in the 65+ age group.



Number of severe complicated influenza confirmed cases by week of onset



* A person who has ILI symptoms become severely ill (includes pulmonary complication, neurologic complication, myocarditis, invasive bacterial infection, or pericarditis) that requires intensive care or results in death within 14 days and with influenza virus infection confirmed by the laboratory is defined as a confirmed severe complicated influenza case.

Number and incidence of severe complicated influenza confirmed cases and deaths by age groups October 1, 2017 to February 5, 2018

Age Group	Cases	Deaths	Cumulative incidence per ten thousand population	Cumulative mortality per ten thousand population
< 3 y	5	0	0.8	0.0
3-6 y	7	0	0.8	0.0
7-18 y	10	0	0.4	0.0
19-24 y	1	0	0.1	0.0
25-49 y	33	6	0.4	0.1
50-64 y	68	13	1.3	0.3
65 +	190	30	6.0	0.9
Total	314	49	1.3	0.2


July 1, 2017 to February 5, 2018 (by flu season)

Age Group	Cases	Deaths	Cumulative incidence per ten thousand population	Cumulative mortality per ten thousand population
< 3 y	15	0	2.4	0.0
3-6 y	16	1	1.9	0.1
7-18 y	13	0	0.5	0.0
19-24 y	7	0	0.4	0.0
25-49 y	78	15	0.9	0.2
50-64 y	157	30	3.0	0.6
65 +	603	102	18.9	3.2
Total	889	148	3.8	0.6



Pneumonia and Influenza (P&I) Mortality Surveillance

Based on the Internet System for Death Reporting (ISDR) surveillance data, the number of deaths attributed to pneumonia and influenza (P&I) during week 4, 2018, was lower than the previous week. The proportion of deaths attributed to P&I for adults aged 65 years and above was the highest among the three age groups (0–49, 50–64, and 65+). Weekly P&I data are available at: <http://nidss.cdc.gov.tw/>.



* Medical institutions are required to report any mortality case to the Ministry of Health and Welfare (MOHW) within 7 days after a death certificate is issued through the Internet System for Death Reporting (ISDR). Either the immediate cause of death or the underlying cause of death was used to identify P&I death cases. Only those with keyword texts containing 'pneumonia', 'influenza' or 'common cold' were counted as a P&I death.

