



## Summary : Week 15 (Apr. 8– Apr. 14, 2012)

Influenza activity in Taiwan peaked at week 2. Seasonal influenza virus type AH3 is predominant in the community.

- Of the 197 specimens tested during week 13, 2012, 39 (19.8%) were positive for influenza viruses, 27 (13.7%) were positive for influenza AH3 virus, 9 (4.6%) were positive for influenza B virus, 3 (1.5%) were positive for pandemic (H1N1) 2009 virus.
- There were 13 new complicated influenza infections and 9 reports of death from complicated influenza infection during week 15. Since July 1, 2011, 1387 complicated influenza infections have been confirmed, including 1025 cases from influenza B virus infection, 276 cases from influenza A virus infection, 65 cases from pandemic (H1N1) 2009 virus infection and 21 cases due to un-typed influenza A virus infection. 466 cases of complicated influenza infections received the intensive care. Since July 1, 2011, there have been 110 reports of death from complicated influenza infection; 88 from the seasonal influenza B strain, 17 from the seasonal H3N2 strain, 3 from pandemic (H1N1) 2009 virus infection and 2 from the un-typed influenza A virus, respectively.
- The number of deaths related to pneumonia and influenza during week 14 was 258, which was above the number of the previous week (254 deaths).
- The rate of outpatient visits for influenza-like illness is 1.22%, a 10.3% decrease compared with the rate of previous week (1.36%).
- The rate of emergency room visits for ILI is 11.42%, a 8.4% decrease compared with the rate of previous week (12.47%).

## Virologic surveillance


The results of tests performed during week 13 are summarized in the table below.

	Data for week 13	Cumulative data since 7/1/2011
<b>Number of specimens tested</b>	197	13693
<b>Number of positive specimens (%)</b>	39(20)	3019(22)
<b>Positive specimens by type/subtype (%)</b>		
<b>Influenza A (% of all positive specimens)</b>	30(77)	509(17)
<b>A (pandemic [H1N1] 2009) (% of all Influenza A)</b>	3(10)	97(19)
<b>A (H3)</b>	27(90)	408(80)
<b>A (H1)</b>	0(0)	0(0)
<b>A (unable to subtype)</b>	0(0)	4(1)
<b>A (subtyping not performed)</b>	0(0)	0(0)
<b>Influenza B</b>	9(23)	2510(83)



**Antigenic characterization:** Taiwan CDC has antigenically characterized 385 human influenza viruses [4 pandemic (H1N1) 2009, 55 influenza A (H3), 326 influenza B] since July 1, 2011. 92.7% (n=51) of the influenza A (H3N2) viruses tested were related to the A (H3N2) vaccine component (A/Perth/16/2009). 13.5% (n=44) of the influenza B viruses tested belonged to the B/Victoria lineage and were related to the B component of the 2011-12 influenza vaccine (B/Brisbane/60/2008).

### Influenza positive tests reported to Taiwan CDC by contracted laboratories, 2010–2012



**Antiviral resistance:** Since July 1, 2011, 28 pandemic (H1N1) 2009, 109 influenza A (H3N2) and 260 influenza B viruses have been tested for resistance to the neuraminidase inhibitors (oseltamivir). The results of antiviral resistance testing performed on these viruses are summarized in the table below.


	Isolates tested (n)	Resistance Viruses, n (%)
		Oseltamivir
<b>A (pandemic [H1N1] 2009)</b>	28	0(0)
<b>A (H3)</b>	109	0 (0)
<b>B</b>	260	0 (0)






## Pneumonia and influenza (P&I) mortality surveillance

During week 14, 2012, 258 deaths reported through the National Death Certificate System mentioned P&I as the cause of death. This number was above the number of previous week (254 deaths). The number of deaths related to P&I for age groups 0–49, 50–64, and greater than 65 years was the highest for adult greater than 65 years of age.

### National pneumonia and influenza mortality Week ending 07/04/2012




## National pneumonia and influenza mortality by age group Week ending 07/04/2012



## Reports of complicated influenza\*


There were 13 new complicated influenza infections (including 6 seasonal influenza AH3 cases, 3 seasonal influenza B cases, 3 cases from pandemic (H1N1) 2009 virus infection and 1 case from un-typed influenza A virus infection) during week 15. Since July 1, 2011, 1387 complicated influenza infections have been confirmed, including 466 cases requiring intensive care. Since July 1, 2011, there have been 110 reports of death from complicated influenza infection; 88 from the seasonal influenza B strain, 17 from the seasonal H3N2 strain, 3 from pandemic (H1N1) 2009 virus and 2 from the un-typed influenza A virus, respectively.

### Number of complicated influenza reports by week of onset 27/6/2010 to present



\*Defined as influenza infection with pulmonary complication, neurologic complication, myocarditis, pericarditis, invasive bacterial infection requiring hospitalization, intensive care or resulting in deaths.

### Rate of complicated influenza reports by age groups 1/7/2011 to present



\*Numbers represent number of complicated influenza reports for that specific age stratum.



## Outpatient and emergency room influenza-like illness surveillance

Nationwide during week 15, 1.22% of outpatient visits reported through the National Health Insurance Database were due to influenza-like illness (ILI), a 10.3% decrease compared with the rate of previous week (1.36%).

The rate of emergency room visits for ILI is 11.42%, a 8.4% decrease compared with the rate of previous week (12.47%).

### Rate of outpatient and emergency room (ER) visits for influenza-like illness

27/6/2010 to present

