計畫編號：DOH98-DC-2037，DOH99-DC-2029

行政院衛生署疾病管制局 98-99 年度科技研究發展計畫

登革熱病媒蚊誘蚊產卵器監測研究計畫

研 究 報 告

執行機構：研究檢驗中心

計畫主持人：吳和生、劉定萍、林立人、鄧華真、張念台

研究人員：吳懷慧、溫在弘、吳智文、翁芝琦、林巧、呂良振、王智源、簡淑婉

執行期間： 98 年 03 月 01 日至 99 年 12 月 31 日

＊本研究報告僅供參考，不代表本署意見，如對外研究成果應事先徵求本署同意

目　　錄

頁　碼

封面

目錄

一、摘要 （ 3 ）

二、前言 （ 5 ）

三、材料與方法 （ 7 ）

四、結果 （ 12 ）

五、討論 （ 47 ）

六、結論與建議 （ 51 ）

七、計畫重要研究成果及具體建議 （ 53 ）

八、參考文獻 （ 55 ）

九、圖 （ 58 ）

十、表 （ 78 ）

十一、附錄一(2010 登革熱防治國際論壇摘要) （ 89 ）

十二、附錄二(文章初稿) （ 96 ）

十三、附錄三(斑蚊空間分布圖) （ 123)

共(125)頁

摘要

本計畫係評估配合人口密度，利用新設計誘蚊產卵器監控台灣登革熱高風險地區之病媒蚊，預估登革熱發生風險的可行性，並於高雄市前鎮區及高雄縣鳳山市配合使用誘蚊機及吸蟲機進行成蚊調查並縣市衛生局所進行之常規監測病媒蚊指數，以比較各種採集方法監測斑蚊之有效性。誘蚊產卵器同時放置產卵棒及塑膠黏片，會顯著降低產卵棒卵數及誘蚊產卵器陽性率，但顯著增加塑膠黏片上之斑蚊成蚊數及誘蚊產卵器陽性率。各種採集方法（誘蚊產卵器、BG-sentinel誘蚊器、背負式吸蟲機、傳統病媒蚊幼蟲指數）均呈現相同之族群消長趨勢，並與一至三個月之登革熱病例數呈現正相關。經過人蚊時空互動分析發現，在疫情的前、中、後期，處於相同的環境因素而感染的疫情傳播模式皆會存在，而因為人的移動造成感染的傳播模式，則主要會出現在疫情的後期。所以我們的結論為(1)這種內置塑膠黏片及產卵棒之小型誘蚊產卵器配合人口密度的放置及衛星定位系統，可同時監測卵、幼蟲、成蟲期的斑蚊密度，亦可預測登革熱發生的風險區塊。(2)誘蚊產卵器、背負式吸蟲機及傳統幼蟲指數均可偵測病媒蚊族群消長，且與一至三月前之病例數呈顯著正相關，建議病媒蚊防治應提早一至三月進行。

中文關鍵詞：登革熱病媒蚊、蚊蟲採集、誘蚊產卵器、空間分析

Abstract

This project is using a population-stratified modified ovitraps to monitor population density of dengue vectors and to evaluate the predictability of dengue epidemic risk in dengue-risk areas of Taiwan. Additionally, BG-sentinel traps and backpack aspirators as well as traditional indices (house index, Breteaux index, and adult index) were also carried to compare the effectiveness to monitor *Aedes* mosquitoes. The wood paddles in the ovitraps with sticky plastics inside significantly attracted more adults and had higher positive rates of ovitraps on the sticky plastics than alone, while the sticky plastics in the ovitraps with wood paddles inside significantly catched more eggs and had higher positive rates of ovitraps than alone. All collection methods showed the same pattern of population succession and had significantly positive association with 1 to 3 month ahead of dengue case numbers. After spatial and time analysis by mosquito and human cases, transmission by mosquioes were found in early, middle, or late phase of outbreak but transmission by human movement only found in late phase. Our conclusion were1)These population-stratified small ovitraps inside with wood paddles and sticky plastics combined with GPS can monitor egg, larvae, and adults of dengue vectors simultaneously and predict dengue outbreak risk areas, and（2) all collection methods (ovitraps, backpack aspirators, traditional indices) can detect the population fluctuation and showed the positive associaton with 1 to 3 months ahead of dengue case numbers. Thus, mosquito control campaign should be launched ahead of occurrence of dengue cases.

Keywords：Dengue vectors, mosquito collection, ovitraps, spatial analysis

前言

登革熱為全球最嚴重之蚊蟲傳播病毒性傳染病，由埃及斑蚊及白線斑蚊傳播。根據世界衛生組織預估每年有五千萬人感染登革熱，其中感染登革出血熱需住院人口約五十萬人，流行地區包括非洲、美洲、東地中海、東南亞及西太平洋，超過100個國家，且區域擴散中，例如2007年在委內瑞拉發生超過八萬個病例流行，其中六千例為登革出血熱病例（WHO, 2010）。台灣登革熱在光復前曾有全島大流行後，沉寂40年後，於1981年在琉球再現(第二型)，1988-1989年在台灣南部引起第一型大流行，確定病例數為4,916例。1990至2003年，其流行幅度為每三至五年小規模流行，每十年大規模流行，至2004年後，每年有小規模流行，流行幅度介於202個病例至2000個病例，流行病毒型別趨於多元。今年截至11月13日止，病例數已超過1000例，流行型別四型，發生於高雄市例（II及III型)、台南市例（IV型)、高雄縣例、台南縣例(I型)、台北縣15例及台北市2例（Taiwan CDC, 2010)。

台灣地區自76年10月於南部爆發登革熱以後，建立登革熱病媒蚊幼蚊調查系統，每個縣市每月調查2個村里，鑑定每隻幼蟲種類，計算埃及斑蚊與白線斑蚊指數、包括住宅指數、布氏指數、容器指數及幼蟲指數。於86年10月為增加調查村里數，以現場目視法鑑定斑蚊幼蟲，亦因目視法無法區分埃及斑蚊與白線斑蚊幼蟲及蛹，所以合併兩種斑蚊種類指數為單一斑蚊指數，調查村里數因而大幅提升。於91年登革熱大流行後，有感於幼蟲密度不能代表登革熱發生的風險，於92年在南部地區新增成蚊調查，以成蚊指數表示，並要求鑑定成蚊種類，了解各地區埃及斑蚊的分布現況。依據最新（97年）登革熱防治工作手冊規定「台南縣、台南市、高雄縣、高雄市及屏東縣每

月病媒蚊密度調查村里數，至少為轄區內總村里數的 50%，其他縣市 6 至 11 月，每月病媒蚊密度調查數，至少為轄區內總村里數的 15%；12 至 5 月，每月病媒蚊密度調查數，至少為轄區內總村里數的 10%」。

登革熱疫苗研發雖然已有曙光，但在未來 5-10 年仍然需要仰賴病媒蚊防治，以控制疫情。所以如何有效監測病媒蚊密度，預估登革熱發生風險，為台灣地區當務之急。新加坡面積 682.2 平方公里，於 2000 年開始放置 2,000 個誘蟲產卵器（30%戶內，70%戶外），每週蒐集 2 片 paddles，計算上面的卵粒，結合地理資訊系統，分析尋找登革熱發生熱區及危險地區（埃及斑蚊高密度區）(Ooi et al. 2006)。香港面積 1,104 平方公里，亦於 2000 年開始，於 2003 年增加至 38 個地點，利用誘蚊產卵器監測轄區白線斑蚊族群，每個點（~0.55 平方公里）放置 55 個誘蚊產卵器，放置約 2,090 個（香港食物環境衛生署網站)。誘蚊產卵器所紀錄的是雌蚊產卵的活動，而與登革熱發生最有關的時期為雌蚊。目前成蚊監測較有效的辦法為背負式吸蟲機、粘紙誘卵器及 BG-Sentinel 誘蚊器（Morrison et al. 2008），而這些方法採集不同蚊蟲族群，例如背負式吸蟲機採集成蚊棲息族群，粘紙誘卵器採集產卵的雌蚊族群，而 BG-Sentinel 誘蚊器採集混合不同的誘引劑，採集吸血雌蚊族群。截至目前為止，雖然有實驗證實經卵傳播的可能性，但在自然界中，仍然由雌蚊吸病毒血後，病毒複製再經過唾液，傳給另外一人。所以吸血一次以上的雌蚊，傳播登革熱的風險就更高，所以配合人口密度監測產卵雌蚊族群應可提供登革熱風險更好的監測指標。本計畫的目的係評估配合人口密度，利用新設計誘蚊產卵器監控台灣登革熱高風險地區之病媒蚊，預估登革熱發生風險的可行性，並於高雄市前鎮區及高雄縣鳳山市配合使用誘蚊

機及吸蟲機進行成蚊調查並縣市衛生局所進行之常規監測病媒蚊指數，以比較各種採集方法監測斑蚊之有效性。

材料與方法

研究地點

選擇過去登革熱常發生的縣市（高雄縣及高雄市）做為此次研究地點。再依據 97 年各鄉鎮市區人口密度及 93 年至 96 年埃及斑蚊分布(埃及斑蚊成蚊占斑蚊百分比),選取人口密度每平方公里超過1500人的鄉鎮市區及埃及斑蚊分布超過 20% 以上，包括高雄市 11 區（新興區、苓雅區、鹽埕區、旗津區、前金區、三民區、前鎮區、鼓山區、左營區、楠梓區、及小港區）及高雄縣 7 個鄉鎮市(鳳山市、梓官鄉、林園鄉、茄萣鄉、仁武鄉、大寮鄉及鳥松鄉)。但因茄萣鄉為單獨的一區，所以不列入此次研究範圍。

登革熱病媒蚊調查方法

此計畫使用之調查方法,包括誘蚊產卵器、BG-sentinel 誘蚊器、吸蟲機等。誘蚊產卵器是一個之細小黑色容器（下底直徑 6.5cm，開口直徑 9 cm x 高 10-13.5cm)，容器裝水 60%，內放置一褐色產卵棒、粘紙(塑膠片)或兩者，並放昆蟲生長調節劑(防止蚊蟲羽化為成蟲)，上面有一個灰色傘頂蓋，外面以白色書寫「疾病管制局實驗進行中請勿干擾」字樣。誘蚊產卵器地點，因埃及斑蚊棲息場所主要在戶內（70-80%)，原則上戶內放置 1 個，戶外 1 個。在放置前，檢查週遭環境，清除積水容器，以免影響結果。誘蚊產卵器放置地點，戶內放置於客廳，戶外放置於隱敝處。另外於大規模實驗前在實驗室進行產卵棒及黏紙效果，並利用高雄世運場點進行前試驗，共放置 22 個點，每個點戶內及戶外共放置 6 個誘卵器，內放產卵棒、黏紙及兩者。

於 98 年 6 月 3-8 日，按照高雄縣及高雄市 97 年度各村里人口密度，以村里為單位，原則上每 400 人放置 2 個誘蚊產卵器，1 個戶內，1 個戶外，共選擇高雄市 3,595 家戶（7,190 個誘蚊產卵器），高雄縣 1585 家戶（3,170 個誘蚊產卵器），合計 10,360 個（圖一及表一），放置期間為 1 年（98 年 6 月至 99 年 6 月）。另外以分層隨機抽樣設計（stratified random sampling）分配三個處理（單獨放置產卵棒、單獨放置塑膠黏片及兩者都放），因埃及斑蚊分布比率會影響處理，再將鄉鎮市區分 2 個區塊，第一個區塊（埃及斑蚊分布介於 82.9%-94.9%以上）包括高雄市苓雅區、旗津區、新興區、鹽埕區、三民區、前金區、前鎮區、楠梓區，第二個區塊（埃及斑蚊分布介於 25.2%-73.9%）包括高雄市左營區、鼓山區、小港區、高雄縣鳳山市、梓官鄉、仁武鄉、林園鄉、鳥松鄉及大寮鄉。

每週取回的產卵棒，計算蚊蟲卵。蚊蟲卵計數完畢，放在塑膠盆孵化，飼養至第 3-4 齡，並在顯微鏡鑑定種類。因為以人工計算卵數，需要花很多的人力，常無法推展，所以發展以影像處理軟體掃描蚊蟲卵（Mello et al. 2008）計算卵數的方法，以代替人力。利用本局建構登革熱病媒蚊調查系統，由高雄市及高雄縣衛生局進行轄區村里調查。依據最新（97 年）登革熱防治工作手冊規定「高雄縣、高雄市及屏東縣每月病媒蚊密度調查村里數，至少為轄區內總村里數的 50%」。調查指數為布氏指數（每 100 戶住宅孳生斑蚊容器數）、住宅指數（每 100 戶住宅孳生斑蚊戶數）及成蟲指數（調查 50-100 戶住宅，每戶平均採集的斑蚊雌蚊數）。另外於高雄縣鳳山市及前鎮區每月隨機各選 2 里，每里選 2 戶，每戶放置 2 個 BG-sentinel 誘蚊器（戶內與戶外各 1 個），放置 1 天（9:00-17:00），並於該里利用背負式吸蟲機吸取成蚊，共 10~15 戶。

統計分析

本研究建立登革熱病媒蚊監測GIS查詢系統，利用統計空間分析軟體分析預估登革熱高風險區，以機率密度得Z值，Z值越高，則代表其涵蓋蚊卵數分佈密度也越高。產卵棒採集之卵數評估病媒分佈的指標。以核心密度估計法(Kernel Density Estimation)計算其在研究區域之密度分佈。核心密度估計為一種可估計隨機變數的機率密度分佈函數的統計方法，其不以行政區域為分析的單位，而是將研究區域切割成緻密之小方格，估計每一個小方格內蚊卵分布的狀況。處理(單獨放置產卵棒、單獨放置塑膠黏片及兩者都放）之間的比較，則使用變方分析（Analysis of Variance, ANOVA）及最小平方差方法（Least Square Difference）分析。因為傳統指數及誘蚊產卵器誘集之斑蚊數低，為比較及統一，所有計算方式均改採陽性率（誘蚊產卵器指數及背負式吸蟲機指數)，有關各種採集方法及其與病例數之相關性，則利用 Pearson correlation 來檢測各種採集方法所採集之登革熱病媒蚊斑蚊族群的相關性。另外使用 paired t 檢測法比較不同採集方法所誘集之雌蚊與雄蚊差異。

病媒蚊體內病毒檢測

1.病媒蚊體內病毒RNA萃取

(1)將由誘卵器及吸蚊機冬季捕獲之蚊子單隻蚊子放入1.5 ml微量試管中，加入0.5ml BA-1溶液，並放入1顆滅菌過的3 mm玻璃珠。

BA-1溶液 1 X medium 199 with Hanks' balanced salt solution ,0.05 M Tris Buffer(PH7.6)

1 % bovine serum albumin

0.35 g sodium bicarbonate /L

100 U streptomycin /L

100 U penicillin

25 ug amphotericin B(Fungizone)/ml

(2)以tissue lyser震盪1分鐘打碎蚊蟲細胞組織。

(3)將均質液，以14000rpm離心10分鐘除去懸浮固體。

(4)取100μl上清液至新的1.5 ml微量離心管中，並加入150μl BA-1溶液，混和均勻。

(5)吸取560μl含有carrier RNA的AVL溶液至1.5ml微量離心管中，並加入140μl步驟4的液體，vortex 1分鐘混合均勻。

(6)室溫(15 ~ 25℃)下作用10分鐘。

(7)加入純酒精560μl，震盪約一分鐘以終止反應。

(8)利用小烏龜離心機離心數秒，將蓋子上的殘留液離下。

(9)將上述混合液630μl分兩次加至QIAamp spin column (放置於2ml collection tube上)，蓋上蓋子，以14000 rpm轉速離心2分鐘，將QIAamp spin column放置新的2ml collection tube上。

(10)小心打開QIAamp spin column的蓋子，加入500μl AW 1溶液，蓋上蓋子，以14000rpm轉速離心2分鐘，將QIAamp spin column放置新的2ml collection tube上。

(11)小心打開QIAamp spin column的蓋子，加入500μl AW 2溶液，蓋上蓋子，以14000 rpm轉速離心2分鐘，倒去下層液。

(12)將QIAamp spin column放置新的1.5ml 微量離心管上，以14000rpm轉速離心3分鐘後，開蓋放置室溫中5分鐘除去多餘的酒精。

(13)將QIAamp spin column放置新的1.5ml 微量離心管上，加入AVE 70μl溶液，靜置於室溫下10分鐘，以14000rpm轉速離心2分鐘。

(14)保存於 -20℃或 -80℃，進行後續病毒檢測用。

2.蚊蟲體內帶病毒SYBR Green real-time RT-PCR檢驗

(Qiagen QuantiTect SYBR Green RT-PCR Kit (Cat NO.204243))

(1)取出Qiagen QuantiTect SYBR Green RT-PCR （Master Mix, RNase-free waer），primers，待溶解後離心（spin-down），立即置於冰上。

(2)依序加入以下試劑

Component	Volumn/reaction	Final concentraction
1. 2x QuantiTect SYBR Green RT-PCR Maser Mix	25μl	1X
2.RNase-free water	Variable	
3.Foeward primer	Variable	
4.Reverse primer	Variable	
QuantiTect RT Mix	0.5μl	
Total	40μl	

配製時多加一份的量，全程於冰上操作，首先加入RNase-free water，再依序加入primer及2x QuantiTect SYBR Green RT-PCR Maser Mix，最後加入QuantiTect RT Mix。

測 Flavivirus group專用primer

FL-F 5’ GCC ATA TGG TAC ATG TGG CTG GGA GC-3’

FL-R1 GTG/T ATT CTT GTG TCC CAT/A CCG GCT GTG TCA TC-3’

FL-R2 GTG ATG CGG/A GTG TCC CAG CCA/G GCT/G GTG TCA TC-3’

(3)加入 40μl Master Mix 至 Q-PCR專用試管

(4)加入檢體、陽性、陰性對組RNA

蚊子檢體：純化出之RNA，取10μl

陽性對照組：以日本腦炎（家蚊屬蚊蟲）或登革熱（斑蚊屬蚊蟲）(黃病毒屬)及Sindy(α病毒屬)病毒RNA稀釋10倍後，加10μl。

陰性對照組：NTC(Non-Template Control)，加10 μl RNase-free water。

(5)蓋上Q-PCR專用試管蓋子，輕輕搖晃混和均勻，離心(spin down)。

(6)用MX4000進行反應及分析結果。

溫度程式 Step	Cycles	Time	Temperature
Reverse Transcription	1	30 min	50℃
PCR Initial Activation	1	15 min	95℃
Denaturation	45	15 sec	94℃
Annealing	45	30 sec	55℃
Extension	45 45	20 sec 30 sec	72℃ 77℃
Dissociation Curve	1 45	1 min	95℃ ↑0.5℃/cycle/ 30sec 68℃

(7)以SYBR Green real-time RT-PCR檢驗結果若Ct值低於40及/或Tm值高於79，則以電泳跑膠比對產物大小。若產物大小與陽性對照組相同，則將產物定序後，進入NCBI資料庫比對。

結果

誘蚊產卵器實驗室試驗

在進行大規模放置前，在實驗室進行產卵棒及黏紙的效果，實驗結果如表二。結果顯示產卵棒與黏紙的效果會嚴重互相影響，且兩者

效果都很好，單純產卵棒的產卵量為 59-345 粒卵，黏紙的黏蚊效果不管是雌蚊或雄蚊均可達 40-60%，產卵量減少至 0-38 粒卵。所以田間實驗同時進行誘蚊牽卵器內單獨放置產卵棒、誘蚊產卵器單獨放置黏紙，及誘蚊產卵器同時放置產卵棒與黏紙等三種試驗。

誘蚊產卵器野外前置試驗

於世運期間放置誘蚊產卵器進行前試驗，共放置 132 個誘蚊產卵器，結果顯示產卵棒每週平均陽性率介於 6.8-18.2%，其中戶內沒有誘集到卵，而戶外陽性率為 13.6-36.4%（表三）。塑膠黏片每週平均陽性率介於 6.8-15.6%，其中戶內為 0.0-4.5%，戶外為 13.6-31.2%。

誘蚊產卵器之誘蚊效果及野外族群消長趨勢

在誘蚊產卵器內放置產卵棒及塑膠黏片，互有影響（表四）。放置塑膠黏片會顯著（$P<0.05$）降低產卵棒誘集之卵數及陽性率，反之，放置產卵棒會增加誘蚊器捕獲之斑蚊數及陽性率。每個誘蚊產卵器同時放置產卵棒及塑膠黏片，產卵棒平均 0.69±0.65 粒卵，陽性率為 2.63%±2.24%，單獨放置產卵棒之平均卵粒為 0.08±0.09 粒卵，陽性率 0.50%±0.48%。放置產卵棒會顯著（$P<0.05$）增加塑膠黏片誘集之斑蚊。每個誘蚊產卵器同時放置產卵棒及塑膠黏片，塑膠黏片誘集之平均斑蚊數為 2.95±1.36 隻，陽性率為 2.28%±1.27%，單獨放置塑膠黏片之平均斑蚊數為平均 1.90±2.50 隻，陽性率 1.63%±1.71%。

就野外族群週消長趨勢分析，埃及斑蚊雌蚊在第 24 開始增加，至 27 週下降，34 週至 42 週呈現浮動，至 43 週至 47 週往下降，48 至 50 週微升後下降，在次年第一週至 16 週均呈現低密度，第 17 週開始上升浮動（圖二）。全年每週塑膠黏片均可採集到埃及斑蚊雌蚊，代表全年埃及斑蚊均有產卵的行為，亦即吸血行為。白線斑蚊族群則無明顯趨勢，僅於第 27 週至 31 週、第 40 週、第 43 至 44 週有採集

到相當數量之雌蚊。對於傳統指數（布氏指數、住宅指數及成蚊指數）而言，與埃及斑蚊雌蚊呈現相同之消長趨勢，於第 24 週開始上升，至 27 週下降後至第 50 週浮動，第 51 週至下降至次年第二週至第 15 週，維持低密度，至第 16 週開始上升。此時雨量及溫度也有明顯的上升變化，氣候對於斑蚊的密度可能有很大的相關性。誘蚊產卵器陽性率之趨勢亦同其他指數（圖三 A），但戶外誘蚊產卵器陽性率於高密度（第 24 週至 39 週）時明顯增加，在低密度時戶內誘蚊產卵器陽性率與戶愴陽性率相同。全年每週產卵棒均可採集到卵，代表全年登革熱病媒蚊均有產卵的行為。

以縣市分別來看，高雄市置有產卵棒(處理 1 與處理 3)的誘蚊產卵器計 4,810 個，高雄縣則有 2,096 個。本研究於所選住戶每戶室內與室外均各置放誘蚊產卵器一個，誘集結果高雄縣市病媒蚊密度高的 6 月~11 月都以室內 ovitrap 陽性率較高(圖三 B 及 C)。此應與夏季高溫，病媒斑蚊喜藏匿棲息較陰涼的室內有關。對於病媒蚊的監測，顯然本 ovitrap 以置放室內效果為佳。每週由產卵棒所採卵經孵化後，三齡幼蟲鑑定埃及與白線斑蚊。兩種病媒蚊合計顯示高雄市的斑蚊數多於高雄縣(圖四 A)，尤其 6 月至 11 月間高雄市病媒蚊數甚高。由誘得卵所孵化之病媒幼蟲檢定得知，高雄縣市終年埃及斑蚊與白線斑蚊的發生消長如圖四（B 及 C）所示。不但每週斑蚊發生量高雄市高於高雄縣，且高雄市埃及斑蚊的發生高於白線斑蚊，高雄縣則以白線斑蚊發生多於埃及斑蚊。

高雄市每週置有黏紙的誘蚊產卵器(處理 2 及處理 3)計 4,782 個，誘捕的病媒蚊成蟲示如圖五 A，以 8 月初至 11 月底誘捕病媒成蚊較高，而各月份均以室內捕獲之成蚊較室外為多。另外幾乎所有月份，田間捕獲成蚊中埃及斑蚊多佔 60%以上(圖五 B)，此與前述以誘

蚊產卵器收集卵後實驗室孵育之蚊蟲相同，高雄市以埃及斑蚊為主要發生之病媒蚊。捕獲成蚊的誘蚊產卵器中，平均每個產卵器約捕得病媒蚊 1~1.5 隻，以每一百個誘蚊產卵器而言，其內若附加黏紙，每週若平均捕得成蚊 2.5 隻以上，則病媒密度值得注意(圖五 C)。高雄縣每週置有黏紙的誘蚊產卵器則計有 2124 個，誘捕的病媒蚊成蟲示如圖六 A，不同於高雄市其捕獲成蚊數較低，且 6 月至 11 月以戶外誘捕之白線斑蚊為多。相對的誘捕蚊數低時，誘得埃及斑蚊的比率較高(圖六 B)。高雄縣捕獲成蚊的誘蚊產卵器中，平均每個產卵器約捕得病媒蚊 1~2 隻，以每百個附加黏紙的誘蚊產卵器而言，每週平均捕得成蚊超過 2.5 隻者僅有 6 週(圖六 C)。

病媒蚊空間分布

若簡略將 6-12 月定義為登革熱流行季，1-5 月定義為非流行季，從 Kernel Density 密度分佈，我們可以很清楚發現，在流行季相較非流行季，病媒蚊在研究區域中的分佈都比較廣(有顏色的區域面積較大)、密度較高(Z 值較大)。而更進一步縮小範圍來看，在流行季時，高雄市三民區的東半部、高雄市苓雅區東半部以及高雄市前鎮區東北部，為病媒蚊分佈密度最高之處（圖七，詳如附錄三)。即便到了非流行季，病媒蚊密度減少，仍可發現高雄市三民區、高雄市苓雅區有病媒蚊穩定分佈。未來在防治上，無論流行季或非流行季，針對高雄市三民區及苓雅區，都應特別加強孳生源清除以及減少病媒蚊的工作。在高雄縣的部分，鳳山市由於鄰近前述高雄市病媒蚊分佈密度高的區域，再加以從圖中亦可見不論是流行季或非流行季，其病媒蚊穩定存在。另外則是林園鄉的南部，在流行季間，病媒蚊的分佈也是屬於密度較高的區域，同樣建議未來應加強相關的防治工作。

依據誘蚊產卵器一至三月冬天的數據，找出 99 年高雄縣市埃及

斑蚊起始族群的重點區塊，提供 99 年度登革熱預防策略防治依據。放置誘蚊產卵器家戶分數計算：每週每一住戶內外有無埃及斑蚊，若有偵測到則給予一分，每戶每週最多兩分(戶內及戶外各一分)。各別將每一住戶分數進行加總，將總分數分成四個等級並以不同顏色區分繪製於 GIS 圖上，然後再以 Kernel Density 密度分布畫出，冬天（99 年 1 月至 3 月 7 日）誘蚊產卵器埃及斑蚊密度圖（圖八），顏色較鮮紅之區塊，經比對後之村里，共 139 里，如表五，為埃及斑蚊起始族群重點區塊，其中包括高雄市三民區 56 里（安泰里、安邦里、民享里、十美里、德行里、德東里、鼎勝里、鼎強里、鼎力里、鼎金里、鼎泰里、鼎中里、鼎西里、灣愛里、灣子里、灣興里、灣復里、灣成里、灣華里、灣勝里、灣利里、灣中里、本和里、本館里、本元里、本武里、本文里、本安里、本上里、本揚里、寶龍里、寶珠里、寶興里、寶中里、寶玉里、寶泰里、寶安里、寶盛里、寶德里、寶獅里、寶國里、寶慶里、寶業里、寶華里、寶民里、安生里、安吉里、安寧里、安康里、安發里、博惠里、港新里、同德里、德仁里、港北里、港東里)、新興區 25 里(順昌里、文昌里、開平里、興昌里、光耀里、德生里、南港里、中東里、成功里、德政里、德望里、華聲里、永寧里、新江里、黎明里、愛平里、東坡里、漢民里、玉衡里、秋山里、明莊里、大明里、榮治里、仁聲里、蕉園里)、苓雅區 19 里(正道里、正心里、正大里、正文里、正言里、永康里、五福里、民主里、五權里、福南里、奏捷里、安祥里、凱旋里、尚義里、中正里、林安里、城西里、意誠里、衛武里)、左營區 11 里(菜公里、新光里、新下里、埤東里、華豐里、新上里、果貿里、果惠里、果峰里、裕豐里、裕興里)、前鎮區 8 里（瑞華里、瑞昌里、瑞興里、瑞崗里、瑞隆里、瑞東里、瑞和里、瑞平里)、小港區 4 里（大苓里、正苓里、二苓里、

三苓里)、楠梓區 2 里（國昌里、裕昌里)、鹽埕區 1 里（博愛里）及高雄縣鳳山市 13 里（善美里、一甲里、富甲里、鎮南里、埤頂里、美齡里、忠孝里、忠義里、中和里、協和里、新強里、興中里、鎮西里)。

登革熱風險評估

一、病例時空分析

2009 年 7 月起至 2010 年 1 月止，本土病例共計 781 位，其發病週次分佈(流行病學曲線)。如圖九所示，2009 年之疫情由第 42 週起邁入高峰，而在 45 週(11 月)左右達到高峰，往前推算四週(日曆週次為 41-45 週)，病媒蚊在研究區域中的分佈恰達到分佈最廣(有顏色的區域幾乎佔了整個高雄市、鳳山市)、密度最高(Z 值在 20 週達到最大值)。

另一方面，圖十描繪了 2009 年 7 月起至 2010 年 1 月止，本土病例的空間分佈。在空間分佈上，高雄市苓雅區、高雄市前鎮區與高雄縣鳳山市，是病例分佈最密集的區域。也前述在偵測病媒蚊時病媒蚊分佈密度最高的區域；疫情在進入高峰前的四週，病媒蚊的分佈廣度與密度也在高峰。在空間上，病例分佈密集的區域也是病媒蚊分佈的區域，該結果說明病媒蚊與造成疫情進入高峰所扮演之重要角色。

從圖十可見，高雄市苓雅區、高雄市前鎮區與高雄縣鳳山市為病例分佈最密集之處。為進一步探知病例與病例之間時間與空間分佈之關聯，研究者計算每一個病例與其他病例發病相隔日數以及間隔距離，並將發病相隔距離與間隔距離繪製成時空散布圖(spatial-temporal scatter plot)，如圖十一。我們從途中可以發現有四個較明顯時空群聚：

第一時空聚集：相隔距離為 0-1 公里，發病相隔日數為 0-10 天

第二時空聚集：相隔距離為 1-2 公里，發病相隔日數為 0-30 天

第三時空聚集：相隔距離為 2-4 公里，發病相隔日數為 0-30 天

第四時空聚集：相隔距離為 4.5-5.5 公里，發病相隔日數為 0-20 天

從時空聚集的意義來解讀，若發病相隔的距離短、相隔的日數也短，其隱含病例與病例之間較是偏向處於相同的環境因素而感染。然而若發病相隔的距離長、發病相隔的日數也長，其隱含病例與病例之間偏向在不同的環境下感染登革熱，換言之，較可能是因為「人的移動」，移動到不同環境受感染或是傳播疫情。進一步將在這四個時空聚集區中，與其他病例關聯最密集的前 10%病例在空間中標示出，如圖十二所示。

從該圖中可以發現在第一個時空聚集區中，與其他病例關聯度高的病例分佈，都相當的集中，然而在第四個時空聚集區中，與其他病例關聯度高的病例分佈，則相對的分散。再次印證了「發病相隔的距離短、相隔的日數也短，其隱含病例與病例之間較是偏向處於相同的環境因素而感染；發病相隔的距離長、相隔的日數也長，其隱含病例與病例之間偏向是因為人的移動造成的感染」這樣的概念。

若再將第一個時空聚集區與第四個時空聚集區中，其他病例關聯度高的病例發病時間分佈畫出如圖十三。從病例發病的週次分佈我們可以得知，不管是在疫情的前、中、後期，處於相同的環境因素而感染的疫情傳播模式皆會存在，而因為人的移動造成感染的傳播模式，則主要會出現在疫情的後期。而在防治的意義上，當一個縣市有發現登革熱病例後，即應針對整個大環境的孳生源投入介入措施。再者，除了對於登革熱感染者居家週遭的環境噴藥外，更要注意登革熱感染者移動的情形，對感染者施以衛生教育，避免外出移動，必要外出，也要著長袖長褲，避免被病媒蚊叮咬而再次傳播。

二、人—蚊互動分析

本研究將前述之第一與第四個時空聚集區中，與其他病例關聯度高的病例為圓心，建立涵蓋 1 公里半徑的方圓範圍，計算其病例發生前四週，一公里的方圓內所涵蓋的偵測點所平均包含的卵數（圖十四所示）。從圖十四中的第一時空聚集病例所涵蓋的卵數來看，前鎮區與苓雅區附近之病媒蚊密度相對較高，需要投入更多資源進行相關的病媒蚊防治，阻斷因環境因素造成的傳播模式，避免疫情更加惡化。然而從第四時空聚集病例所涵蓋的卵數來看，其病例分佈與卵數高低已無明顯相關，說明了登革熱防治在初期的重要性，一旦疫情發展到後期，由於人的移動複雜及相關的影響因子眾多，屆時要進行相關防治介入措施就會相當困難。

將 98 年度病例分為流行初期（8 月）、流行高峰（11 月）及非流行季（2 月）三個時期（圖十五 A），而流行初期蚊蟲採集之卵數最多，流行高峰次之，非流行季最少（圖十五 B）。依據此三個時期，畫出病媒蚊空間分布圖（圖十六 A）、病媒蚊群聚區塊（圖十六 B）及病媒蚊移動（圖十六 C）。以病媒蚊空間分布乘以人口密度預估登革熱風險區塊（圖十七 A），最後標示預估群聚區塊及 2009 年 8 月至 2010 年 1 月實際發生地區及群聚區塊（圖十七 B）。再將此兩圖再加上病媒蚊空間分布圖做一比較，發現小港區之疫情為流行初期，但因為防治沒有落實，導至病例延至第 52 週（圖十八 A）；三民區則為流行高峰病媒蚊起來後之疫情。2010 年亦有類似狀況，但仍需病例個案資料分析（圖十八 B）。

各種採樣方法之比較及族群月份消長趨勢分析

高雄市前鎮區及高雄縣鳳山市放置誘蚊產卵器位置如圖十九，各種採樣方法所採集之住家型蚊蟲種類，包括熱帶家蚊、埃及斑蚊、白線斑蚊、白腹叢蚊、黃尾家蚊等五種蚊蟲種類，而三種採集方法所捕

獲之蚊蟲種類組成稍有不同(表六)。誘蚊產卵器共捕獲 19,894 隻蚊蟲，以熱帶家蚊最多，共 17,973 隻 (10,246♀7,727♂)，佔 90.3%，其次為埃及斑蚊 1,136 隻(679♀457♂)，佔 5.7%，接著為白線斑蚊 227 隻(189♀38♂)，佔 1.1%，另有 2.6%之斑蚊 517 隻(298♀219♂)因身上之花紋無法辨識，無法完成種類鑑定，其他種類低於 1%者尚包括白腹叢蚊及黃尾家蚊。BG-sentinel 誘蚊器共捕獲 579 隻，以埃及斑蚊 265 隻(170♀95♂)最多，佔 45.8%、熱帶家蚊 207 隻(155♀52♂)次之，佔 35.8%、接著為白腹叢蚊 93 隻(88♀99♂)，佔 16.1%及白線斑蚊 14 隻，佔 2.4%。背負式吸蟲機共捕獲 1,279 隻，以熱帶家蚊 773 隻(457♀316♂)最多，佔 60.4%，埃及斑蚊 342 隻 (173♀169♂)次之，接著為白腹叢蚊 130 隻 (87♀43♂)，佔 10.2%及白線斑蚊 34 隻(26♀8♂)，佔 2.7%。三種採集方法中，僅 BG-sentinel 吸蟲器採集之蚊蟲，雌蚊數顯著 ($t4=4.16$，$P<0.05$) 多於雄蚊數，而其他兩種採集方法，雖採集較多之雌蚊，但沒有顯著性差異 ($P>0.05$)。

將各項調查方法轉換為住宅指數(住宅陽性率)做為比較，結果顯示住宅指數以誘蚊器最高，平均為 51.0% (0-100%)，吸蟲機次之，為 29.0%(0-67.5%)，接著為各縣市衛生局人工調查之住宅指數 3.3% (0.73-5.97%) 及誘蚊產卵器 2.9% (0.53-6.14%)(表七)。各種採樣指數之趨勢變化皆相似於傳統指數 (布氏指數、住宅指數及成蚊指數)，而這些趨勢與溫度及降雨量的趨勢一致，在 2009 年 8 月趨勢開始下降，並在 2010 年 3~4 月開始上升。以上趨勢顯示蚊蟲的密度與溫度及降雨量有關，而本土登革熱病例則延遲兩個月才到達高峰(154 例)，顯示蚊蟲密度的上升後，登革熱病例可能才開始上升(圖二十)。

不同採集方法間的相關性分析如表八所示。而誘蚊產卵器指數、

誘蚊器指數及吸蟲機指數為捕獲斑蚊陽性戶數的比例(住宅指數或住宅陽性率)。布氏指數及住宅指數間有最高的相關性(r=0.997，*P*<0.00)。住宅指數與布氏指數則與誘蚊產卵器指數(r=0.92-0.93，*P*<0.00)及吸蟲機指數(r=0.93-0.94，*P*<0.00)有較高的相關性，與誘蚊器指數則有較低的相關性(r=0.65-0.68，*P*<0.05)，但都具有顯著性的意義存在。誘蚊產卵器指數也與吸蟲機指數有高的相關性(r=0.98，*P*<0.00)，但對於成蚊指數及本土登革熱病例數無關。誘蚊器指數與大部份其它的指數有較低的相關性(r=0.58-0.68, P<0.05)。吸蟲機指數則與大部份的傳統指數及誘蚊產卵器指數有高的相關性(r=0.93-0.98，*P*<0.00)，成蚊指數及本土登革熱病例與其他指數則無相關性(*P*>0.05)，但成蚊指數與吸蟲機指數則有低的相關性(r=0.58，*P*<0.05)。因斑蚊密度的上升趨勢較本土登革病例數還早，因此將本土登革病例數往前推移 1 至 4 個月進行相關性分析，如表九，結果顯示 1 至 3 個月的推移皆與布氏指數、住宅指數、誘蚊產卵器指數及吸蟲機指數有相關性(r=0.56-0.81，*P*<0.05)，其中以往前推移 2 個月的相關性最高(r=0.67-0.81，*P*<0.05)，但與於成蚊指數與誘蚊器指數則無相關性。

冬天斑蚊成蚊登革熱病毒檢驗

為了檢測登革熱病毒是否存在於冬季殘留的斑蚊族群中，將自高雄市前鎮區及高雄縣鳳山市以誘蚊器及吸蟲機捕獲之斑蚊進行登革熱病毒檢驗，共檢測埃及斑蚊 82♀44♂，白線斑蚊 2♀，結果顯示只有 3 隻雌埃及斑蚊檢測出 flavivirus 弱陽性，分別為誘蚊器 1 隻(6 月鳳山市 Ct 值 40,Tm 值 81.65)，吸蟲機 2 隻(4 月鳳山市 Ct 值 40,Tm 值 80.05；6 月前鎮區 Ct 值 40,Tm 值 82.34)，經定序後則與蚊子一種 flavivirus (cell fusing agent virus, CFAV)序列相似度最高，

而與登革熱病毒序列相似度較低，因此在冬季的斑蚊中，並未檢測到登革熱病毒的存在。

自動計算卵數軟體與網站登錄之研發

由於影像判讀程式係透過影像像素(pixel)大小及群集數量的設定來區分蟲卵的數量，故程式設計讀取外部參數檔來取得此類參考值，如此一來，使用者即可自行進行微調，藉以應用於其他物種不同蟲卵大小之判讀，外部參數檔內容將於後說明。

1.作業程序及範例說明

（1）利用影像判讀程式[紙張列印]功能，分別輸入各監測點編號(最多五碼)及日期後，列印出客製化設計之各監測點的蟲卵棒掃瞄底稿 A4 樣本紙一份(共二張)，分別為蟲卵棒**正面**的監測點室內及室外底稿 A4 樣本紙一張、**反面**的監測點室內及室外底稿 A4 樣本紙一張。



影像判讀程式[紙張列印]操作畫面



列印的**正面/反面**監測點室內及室外底稿 A4 樣本紙範例

（2）作業員於外業取得之監測點室內及室外蟲卵棒，分別自行於蟲卵棒上方 1/4 處以上位置註記正反面標記後，分別將**正面**朝上的室內及室外蟲卵棒，依據前述列印之**正面** A4 樣本紙上所標示之室內及室外的位置，將蟲卵棒浮貼於所對應之室內及室外標示位置上，浮貼原則以蟲卵棒底部恰好完全蓋過標示位置為準，浮貼完成後將 A4 紙張放入掃描器，並設定 ***200 DPI*** 的掃瞄器像素設定，取得監測點正面的蟲卵棒掃瞄影像檔。

（3）同步驟二流程製作反面之蟲卵棒掃瞄影像檔。

（4）透過影像判讀程式[圖檔分析]功能，點選{載入圖檔}按鈕開啟影像檔讀取視窗，於視窗內選取步驟二取得之正面的蟲卵棒掃瞄影像檔後，判讀程式將立即進行蟲卵數判讀分析並將監測點正面的室內及室外蟲卵棒數量輸出成 Excel 檔並開啟，此外，程式亦將原始圖檔及分析後圖檔分別顯示於系統操作介面的左右方供使用者參考判讀結果準確性。

（5）同步驟四流程選取反面的蟲卵棒掃瞄影像檔，並取得反面

的監測點室內及室外蟲卵棒數量 Excel 檔。(如圖 5.1-4 所示)。

(6)全部取得同一監測點之室內/室外的正反面蟲卵數資料後，即可彙整記錄供後續使用。

浮貼上**正面/反面**的室內及室外蟲卵棒範例

正面掃瞄影像判讀成果如下↓

Microsoft Excel - 00001200905210.xls

	A	B	C	D	E	F
1	掃描編號	地點編號	日期	正/反面	室內蟲卵數	室外蟲卵數
2	00001-20090521-0	[00001]	20090521	正面	46	146
3						

反面掃瞄影像判讀成果如下↓

	A	B	C	D	E	F
1	掃描編號	地點編號	日期	正/反面	室內蟲卵數	室外蟲卵數
2	00001-20090521-1	[00001]	20090521	反面	65	56
3						

影像判讀程式取得之**正面/反面**室內及室外蟲卵棒數量 Excel 檔範例

2. 影像程式外部參數檔內容說明：

影像程式外部參數檔檔案名稱為[Setup.ini]，其檔案位置需與影像程式同一目錄，其內容共計有五列，如下所示：

[CONF] ……**第一列**

AREAID=00001 ……**第二列（固定五碼）**

THRESHOLD=160 ……**第三列**

OverNum=100 ……**第四列**

PerPointNum=5 ……**第五列**

各列內容說明如下：

（1）第一列內容

代表表頭，無特殊意義請保留。

（2）第二列

[AREAID]代表區域代號，此代號為產製 A4 樣本紙之條碼所用，若有同一監測站號但不同區域者，請修改**[AREAID]**的內容，例如：高雄市、高雄縣所屬的監測站皆有同一編號的測站，則分別執行高雄市、高雄縣的影像判讀程式時，須於**[AREAID]**參數中，分別輸入不同值藉以做區分，高雄市輸入 **AREAID=00001**；高雄

縣則輸入 **AREAID=00002**。

（3）第三列

[THRESHOLD]是演算法中將影像像素轉化成二值化像素（僅有黑與白）的一個參數值，愈大的參數值，二值化後出來的雜訊愈少，但過大卻會把原值轉成白色，變成白白的一片；反之，若參數值愈小的話，則可能把影像值都轉成黑色變成一團黑，故建議使用者非必要請勿修改此參數，請保留程式測試後的最佳值 **THRESHOLD=160**。

（4）第四列

[OverNum]是影像程式判讀黑點像素超過此值時，則將此點數視為無效，不計入蟲卵數，例如蟲卵重疊過度，則程式判讀必然有相當誤差，此時則提醒使用者須進行人工判讀。

（5）第五列

[PerPointNum]是影像程式判讀黑點像素超過此值時，則視為一個蟲卵數，使用者可調整此參數藉以調整影像判讀程式於其他不同種類、不同大小蟲卵數量判讀時之應用。

登革熱病媒蚊監測 GIS 查詢系統

1. 系統開發環境及整合

現階段電腦之發展趨勢走向雲端運算技術，亦即表示未來的應用系統將大量建置於遠端伺服器，透過網路服務將運算分析結果提供給使用者，因此，本計畫嘗試導入雲端運算技術概念，建立一套不需使用者下載任何的外掛程式或安裝工具即可進行操作的系統，環顧目前市面上可採用作為網際網路電子地圖服務之軟體甚多，包括 ArcGIS Server、MapGuide、SuperWebGIS、Skyline 及 Google Map 等，於考量本計畫之需求、系統建置費用、系統整合效率、軟體支援性及以網

頁為操作平台的雲端運算架構，並以使用介面操作方便、簡單的功能設計構想下，故設計之系統架構將於 Windows Server 作業系統內採用 Microsoft Visual Studio .Net 作為網頁系統的開發工具平台，搭配後端資料庫工具軟體 Microsoft SQL Server 與網際網路電子地圖元件(Google Map)，進行「疾病管制局登革熱蟲卵監測資料查詢顯圖」系統之開發，本系統所使用之作業系統、工具軟體下表。

系統開發所使用軟體及費用一覽表

軟體項目	軟體名稱	備註
作業系統	Ms Windows Server	一般隨伺服器一同銷售
資料庫系統	Ms SQL Server 2005	需購買
網頁伺服器	Ms IIS 6.0	Windows Server 內建功能
程式開發語	Ms Visual Studio	僅開發端才需購買(架設
網際網路電子地圖元	Google Map	Google 公司提供個人及學術單位免費使用

2. 系統架構

「登革熱病媒蚊監測 GIS 查詢系統」係以網際網路做為主要的作業平台，採三層式的架構建置，系統運作架構示意如下圖所示。其觀念是將原先二層式的主從架構擴增為三層式(Three tier)的架構，甚至是多層式(multi-tier)的分散架構，將運算邏輯(模式)獨立出來自成單獨的一層，如此一來，資料庫、運算邏輯(模式)與使用者界面各司其職，各自存放在最適宜的電腦上執行，不但資料甚至運算邏輯(模式)也可以共享使用，可達到分散化運算與資源共享的目的，系統的彈性與維護性也更佳。本系統的網頁伺服器及網際網路地理資訊系統置於

第二層(應用伺服器層)，此層主要為應用軟體之功能執行，為接收使用者指令後，執行啟動應用程式，並視需求向第三層之資料庫伺服器，存取所需之資料，此層所耗用之電腦資源最多，故須以較佳之設備執行。本系統資料庫設置於第三層(資料庫伺服器層)，此層主要負責指標調查資料的儲存與管理，並以 Microsoft SQL Server 為資料庫的伺服器，提供模式執行所需的基本資料、執行結果之儲存以及使用者端對於各監測資料查詢的服務。



三層式系統架構示意圖

3. 系統功能架構及操作說明

「登革熱病媒蚊監測 GIS 查詢系統」主要提供使用者透過網頁瀏覽器查詢各項監測調查資料，並搭配 Google Map 空間分佈展示功能，提

供各類資料之空間分佈資訊，此外，亦透過進階的資料統計分析功能，提供各式圖表之統計展示，包括長條圖、歷線圖等，以提供不同需求之使用者使用，系統操作畫面如下圖所示。

登革熱病媒蚊監測 GIS 查詢系統畫面

各操作區群組說明如下：

（1）圖面操作區

圖面操作區採用 Google Map 元件進行設計，因此其操作方式與 Google Earth 非常相似，可讓使用者非常容易上手操作，本系統設計之主要功能，在於提供使用者利用網路瀏覽器查詢登革熱蟲卵監測資料的空間分佈情形，藉由指定顯圖之資料來源及設定展示樣式(資料範圍、顏色、透明度即屬性資料等)，系統即依據使用者設定條件進行空間顯圖展示。同時亦設計提供圖面操作工具及不同主題圖之呈現方式，方便使用者進行圖面之查詢操作，功能說明如下：

A. 圖面範圍為全球範圍。

B. 提供四種主題底圖資料，分別為：一般地圖、衛星地圖、地形地圖及混和地圖(衛星+道路)，如下圖所示。

C. 座標系統為經緯度座標。

D. 圖面操作功能如圖面放大、縮小、平移外，亦提供比例尺及鷹眼圖工具，此外亦提供快速鍵(Double Click、滑鼠滾輪)放大、縮小功能。

E. 不提供顯圖影像下載功能。

一般地圖	衛星地圖
地形地圖	混和地圖

圖面操作區主題圖樣式範例畫面

2. 顯圖設定區

顯圖設定區主要提供使用者針對呈圖內容進行資料項目的篩選、統計後直接於圖面上顯示，系統提供最多十組的不同資料範圍顯示外觀設定，透過各不同資料範圍的展示外觀設定(顏色、透明度等)，可方便使用者於圖面上觀察出資料的空間分佈情形，

此外亦提供使用者點選圖面資料查詢屬性資料功能，本區畫面如下圖所示，各子功能操作方式說明如下：



顯圖設定區範例畫面

（1）【清除圖面】按鈕

按下此按鈕可將系統套疊之蟲卵監測點位置圖資清除，若同時勾選不顯示縣市行政區邊界，則圖面將只顯示原始主題圖，若勾選顯示縣市行政區邊界，則圖面將只顯示原始主題圖與縣市行政區邊界如下圖所示。

原始圖面範例	執行後圖面範例

【清除圖面】按鈕執行前後結果比較畫面

（2）〈疊圖顯示設定〉功能區

此功能區主要提供使用者進行疊圖、展示屬性資料及指定呈圖的資料來源及統計方式，說明如下：

A. 疊圖：

使用者可勾選【顯示高雄市邊界】來決定是否於圖面上顯示高雄市邊界範圍。

B. 展示屬性資料：

使用者可勾選【顯示監測點屬性資料】來決定顯示於圖面上之監測點資料是否要同時展示該監測點的屬性資料，例如測點編號、處理方式、所在室內外統計方式、週數及監測調查值等，如下圖所示。

不啟用屬性資料展示功能	啟用屬性資料展示功能

展示屬性資料功能啟用結果比較畫面

C. 呈圖資料來源：

使用者可指定利用『蟲卵數』或『白線斑蚊數』或『埃及斑蚊數』調查資料來進行篩選統計及分層範圍顯示之資料來源。

D. 資料統計方式：

由於每個監測點各有室內及室外的監測資料，故系統提供使用者可指定利用『室內』或『室外』或『室內＋室外』的統計資料來進行篩選統計及分層範圍顯示。

（3）〈顯圖資料 - 篩選條件設定〉功能區

此功能區主要提供使用者進行指定資料項目的篩選條件設定，可進行篩選之項目包括：指定縣市、指定鄉鎮區、指定統計的年度月份及監測資料的處理方式(產卵棒、黏蚊紙或產卵棒+黏蚊紙)等，在進行篩選條件時，勾選該項目的指定條件，則所屬的選項才會採用，如下圖所示。

顯圖資料-篩選條件設定說明畫面

（4）〈顯圖外觀設定 – 分層分色特徵設定〉功能區

此功能區主要提供使用者篩選統計資料的分層展示外觀設定及執行顯圖功能，使用者可於此功能區自行設定分層的資料統計範圍值(最小值、最大值)及顯示外觀(顏色、透明度)，操作畫面如下圖所示，說明如下：



顯圖外觀設定-分層分色特徵設定說明畫面

E. 指定分層數量：

使用者可指定系統列出可供選擇的分層數量，最多為 10 組。

F. 指定統計資料範圍

此功能區主要提供使用者自行指定各分層顯示數值範圍、顏色及透明度等，在進行分層設定時，**務必勾選該分層設定，則所屬的分層設定值才會採用**。

於所屬分層數值範圍分別輸入起始值及最大值，則系統將針對統計結果屬於本數值範圍之資料進行顯圖功能，資料篩選範圍數值範例如下所示。

起始值<= 統計資料 <最大值

G. 指定顏顯圖色：

於所屬分層顏色設定點選顏色面版即顯示顏色設定面版(如下圖所示)，其操作步驟如下：

a. 點選顏色設定面版。

b. 於顏色設定面版內點選顏色或輸入顏色之 RGB 數值。

c. 點選【OK】完成設定，若未點選，則顏色將不改變。

顏色設定面版操作說明畫面

H. 指定顯圖透明度：

設定所屬分層顯示之透明度，提供五種選取選項，依序為不透明(0%)、25%透明、50%透明、75%透明及全透明(95%)，如圖 5.9-10 所示。



圖 5.9-10. 透明度設定說明畫面

篩選結果資料展示區

使用者按下【執行顯圖】按鈕後，系統即根據使用者所設定之資料來源、篩選條件、統計條件及分層分色特徵設定進行顯圖作業，完成後將自動切換設定資料查詢區之展示頁籤於「GMap 查詢結果展示」分頁，並同時顯示篩選統計資料之結果及資料筆數，如圖 5.9-11 所示。



圖 5.9-11. 顯圖設定篩選結果資料展示畫面

資料查詢區

目前資料查詢區主要以資料表展示方式提供使用者針對原始監測資料、或篩選統計分析後之資料進行查詢，並可將查詢結果下載成 Excel 格式於本機端開啟，目前本區提供二個資料表之查詢頁籤及一個配合顯圖成果之篩選統計資料展示頁籤，各頁籤分項功能說明如下：

原始資料查詢展示

主要提供使用者進行資料庫內原始蟲卵監測資料篩選查詢功能，使用者可透過資料篩選功能設定各類條件，系統即將資料庫內符合該篩選條件之資料以表單方式呈現於網頁上，並同時提供資料筆數顯示、資料表單下載及單一監測點定位顯圖等功能，操作介面如下圖所示，說明如下：

原始資料查詢展示頁籤操作畫面

篩選條件操作區：

使用者可於此操作區分別設定各篩選條件，包括：縣市別、鄉鎮區名、監測分區、監測點編號指定、處理方式、室內外、週別、監測日期及各類監測資料範圍設定等，使用者按下【蟲卵監測資料查詢】按鈕即可執行篩選查詢，**各篩選條件將以聯集(同時符合)的方式執行**。

I. 資料表操作區：

使用者可於資料表點選頁次符號內進行分頁切換功能；點選欄位名稱進行欄位內容資料的順向排序或反向排序；點選單筆記錄的【定位】圖示進行該筆紀錄的定位顯圖功能(如下圖所示)。

單筆蟲卵監測資料定位顯圖展示畫面

J. 資料成果統計顯示區：

主要展示所篩選資料的筆數及提供使用者下載 Excel 檔於本機端開啟。

1. 篩選統計結果展示

此功能區主要功能為提供使用者篩選原始蟲卵監測資料並依據指定項目的進行資料統計，此頁籤內含二個子頁籤，分別為〈篩選統計-條件設定〉及〈篩選統計-成果展示〉，使用者於〈篩選統計-條件設定〉執行篩選成果後，系統將自動切換操作畫面至〈篩選統計-成果展示〉頁籤，各子頁籤功能說明如下：

A. 〈篩選統計-條件設定〉頁籤操作區：

此操作區主要分為三部分(如下圖所示)，分別為「監測值統計項目」設定、「監測值篩選條件」設定及「統計分類項目」設定，各分區說明如下：

篩選統計-條件設定操作畫面

1. 「監測值統計項目」設定

此分區主要提供使用者指定欲進行統計的資料項目及統計值，目前提供之項目有蟲卵調查資料之蟲卵數、白線斑蚊數、埃及斑蚊數及成蚊調查資料之白線斑蚊、埃及斑蚊及其他斑蚊資料等，統計值包括合計值、平均值、最大值、最小值及調查次數等五種，成蚊調查資料並可區分公蚊、母蚊及合計等三種。。

2. 「監測值篩選條件」設定

此分區主要提供使用者進行篩選條件之設定，提供之篩選條件包括所在縣市、所在鄉鎮區、所在監測分區、指定室內外、指定處理方式、指定測站編號及指定週數範圍等，使用者若指定篩選條件，則系統將指針對符合條件之原始監測資料進行統計，若未指定則針對全部的原始資料進行統計，例如使用者若指定測站於「室內」，則系統即針對

所有位置於「室內」的資料進行統計，若未指定，則系統即針對所有位置於「室內」及「室外」的資料進行統計。

3. 「統計分類項目」設定

此分區主要提供使用者進行統計分類項目之設定，提供之項目包括縣市名、鄉鎮區名、監測分區、室內外、處理方式、指定測站編號及指定週數等，使用者若勾選指定的分類項目，則系統將針對該項目進行分類統計，且**分類項目統計結果將以相乘的方式擴增**，例如若只勾選縣市名，則統計的結果將只有二筆紀錄，分別為「高雄市」及「高雄縣」，若同時勾選縣市名及處理方式，則統計的結果將有六筆紀錄，分別為高雄市的「產卵棒」、「黏蚊紙」及「產卵棒+黏蚊紙」與高雄縣的「產卵棒」、「黏蚊紙」及「產卵棒+黏蚊紙」，如下兩圖所示，所有篩選設定於按下按鈕後即可進行統計，完成後將自動切換畫面至〈篩選統計-成果展示〉頁籤並展示篩選統計成果資料表。

只有指定「縣市名」統計分類項目

〈篩選統計-條件設定〉畫面



〈篩選統計-成果展示〉畫面

只指定「縣市名」統計分類項目結果展示畫面

同時指定「縣市名」及「處理方式」統計分類項目

〈篩選統計-條件設定〉畫面

〈篩選統計-成果展示〉畫面

同時指定「縣市名」及「處理方式」統計分類項目結果

B. 〈篩選統計-成果展示〉頁籤操作區：

此頁籤內容顯示與前述原始資料查詢展示頁籤內容類似，其操作亦相同，在此不再重複，畫面如下圖所示。

篩選統計-成果展示畫面

2. GMap 查詢結果展示

此頁籤展示之資料表為配合顯圖篩選統計查詢之成果，操作方式同前述資料表說明。

4. 情境案例範例介紹

情境模擬一：使用者欲查詢所有原始蟲卵監測資料的各鄉鎮區統計，統計項目包括蟲卵數及白線斑蚊歷次調查記錄的平均值、最大值及最小值，統計資料不區分「室內」或「室外」，需合併統計「室內」+「室外」的結果。

設定畫面：

設定畫面於【資料查詢區】-【篩選統計結果展示頁籤】-【篩選統計結果展示頁籤】狀態，並設定如下畫面後按下【執行篩選統計查詢】按鈕。



情境模擬一設定畫面

成果畫面：共計有高雄市 11 個行政區及高雄縣 6 個鄉鎮的統計資料共 17 筆記錄。

註：資料庫內目前僅有高雄縣 6 個鄉鎮的調查資料。

情境模擬一成果彙整畫面

情境模擬二：使用者欲查詢所有原始蟲卵監測資料的各縣市、前 6 週各週的統計，統計項目包括蟲卵數及白線斑蚊歷次調查記錄的平均值、最大值及最小值，統計資料需區分「室內」及「室外」的結果。

設定畫面：

設定畫面於【**資料查詢區**】-【**篩選統計結果展示頁籤**】-【**篩選統計結果展示頁籤**】狀態，並設定如下畫面後按下【執行篩選統計查詢】按鈕。



情境模擬二設定畫面

成果畫面：共計有高雄市、高雄縣、室內及室外各 6 週的統計資料計 24 筆記錄。

分類項目：2 個縣市別*2 個室內外*6 週別 (=24)

縣市名	室內外	週數	蟲卵數平均值	最大蟲卵數	最小蟲卵數	白線賽效數平均值	最大白線賽效數	最小白線賽效數
高雄市	外	5	21	127	1	2	38	0
高雄市	外	6	25	317	1	1	15	0
高雄縣	內	1	15	41	2	1	9	0
高雄縣	內	2	14	41	3	2	14	0
高雄縣	內	3	22	129	0	1	18	0
高雄縣	內	4	31	132	2	3	38	0
高雄縣	內	5	17	83	2	2	12	0
高雄縣	內	6	30	108	5	4	17	0
高雄縣	外	1	34	79	1	2	35	0
高雄縣	外	2	24	54	1	1	17	0

1 2 3

縣市名	室內外	週數	蟲卵數平均值	最大蟲卵數	最小蟲卵數	白線賽效數平均值	最大白線賽效數	最小白線賽效數
高雄縣	外	3	26	95	1	3	35	0
高雄縣	外	4	29	122	3	4	64	0
高雄縣	外	5	27	158	2	5	47	0
高雄縣	外	6	23	147	2	5	27	0

1 2 3

情境模擬二成果彙整畫面

情境模擬三：使用者欲於圖面顯示高雄市第 15 週的蟲卵數調查結果空間分佈狀態，蟲卵數的計算採用室內與室外合計的方式，總共要區分 3 種等級，各等級的蟲卵數合計值範圍如下所示，顏色依序為綠色、藍色、紅色且皆為不透明狀態，圖面需套疊高雄市邊界且不顯示屬性資料。

第一級：1<= 合計蟲卵數<50

第二級：50<= 合計蟲卵數<100

第三級：100<= 合計蟲卵數<1000……..(大等於 100)

設定畫面：

設定畫面於【**顯圖設定區**】設定如下畫面後按下【執行顯圖】按鈕。

情境模擬三設定畫面

成果畫面：



情境模擬三成果畫面

計畫查核

本計畫利用大批志工進行誘蚊產卵器監測，數據之可靠與否端賴工作人員之工作態度、居民配合度等因素息息相關，故於計畫執行期

間透過查核的方式發覺問題，98 年度屏科大已查核前鎮區、苓雅區、鼓山區、三民一區、三民二區、左營區、楠梓區、小港區、旗津區、前金區及高雄縣林園鄉、鳳山市、大寮鄉、梓官鄉，合格率變化很大，戶外為 7.1-100.0%，戶內為 17.3-100.0%（表十）。不合格的原因包括沒有放置、內外容器顛倒、一戶兩個容器均放於戶外（因住戶改變放置意願）、水不夠、黏紙放返等因素。疾病管制局亦不定期進行小規模查核，11 月 05-06 與屏科大共同查核 30 戶，合格率介於 62.5-100.0。99 年度疾病管制局與屏科大共同查核鼓山區、三民區及鳳山市(表十)，合格率戶外為 30-90%，戶內為 10-40%。不合格的原因包括水量過多或過少、容器皆放室內或室外、容器不見等因素，皆已改正缺失。為避免計畫結束後誘蚊產卵器遺留造成滋生源，回收的查核也列為重點項目，疾病管制局與屏科大則共同查核苓雅區、前金區及鳳山市，其室內外合格率皆為 90-100%(表十)。

成本效益分析

本計畫所需誘蚊產卵器監測放置1個誘蚊產卵器監測1日費用為79.27元，監測53週為1,382.91元，放置10,360個誘蚊產卵器費用為14,326,947元（表十一）。另估算登革熱緊急防治一個病例（防治半徑五十公尺範圍）共需12,000-24,000元。若能利用此次誘蚊產卵器監測所得之埃及斑蚊起始族群地區及登革熱高風險地區進行登革熱預防策略，藉降低埃及斑蚊密度，降低病例流行幅度，至少減少596-1190例，進而防止登革出血熱之流行，此研究是值得鼓勵。

討論

本計畫所使用的採集方法（誘蚊產卵器、BG-setntinel 吸蚊器，及背負式吸蟲機）及傳統病媒蚊幼蟲指數（布氏指數及住宅指數），皆可有效偵測斑蚊密度消長，其中誘蚊產卵器配合衛星定位及人口密

度放置，可有效分析病媒蚊時空分布，找出埃及斑蚊每年起始族群的區塊，並藉人蚊互動模式闡述疫情中的兩種傳播模式—當地的環境因素傳播與人口移動的兩種不同的潛在傳播模式，不管是在疫情的前、中、後期，處於相同的環境因素而感染的疫情傳播模式皆會存在，而因為人的移動造成感染的傳播模式，則主要會出現在疫情的後期。所以對於病例的居家週遭噴藥，僅能防治因當地的環境因素感染的傳播模式，對於人口移動造成的傳播模式並無改善的效果。

主要由埃及斑蚊與白線斑蚊傳播的登革熱(Dengue fever)與登革出血熱(dengue haemorrhagic fever)為一種藉蚊蟲傳播感染人類之急性病毒病，在東南亞、西太平洋、拉丁美洲與加勒比海地區的許多國家，每年全世界估計有 5 千萬個病例，是相當重要的公衛議題(Farrar, et al. 2007)。Lloyd (2003) 對於美國預防與防治登革熱最佳實務的報告中，建議了十項必要的工作，其中第一項就是建議整合流行病學與昆蟲學的監測調查(Integrated epidemiological and entomological surveillance)。當然，除了要有正確快速的臨床病毒偵測外，更要有病媒蚊發生與密度的可信監測，才能整合成有用的資料，作為疫病流行情況的處理或防治決策擬定的參考。本計畫係本省自 1987 年登革熱暴發以來首次以人口數為基準，於高雄縣市進行之大規模置放新設計之誘蚊產卵筒(ovitrap)，進行斥堠式(scouting)病媒蚊的監測。探討此種病媒監測方法實際之可行性與其對病例發生間的關聯性。

以誘蚊產卵器誘集成蚊產卵作為埃及斑蚊調查的工具是 Fay and Pery (1965)在實驗室發展出，之後 Fay and Eliason (1966)再加改良而成。這種標準的 CDC ovitrap 誘蚊產卵器是將高 13 cm、直徑 6cm 的廣口瓶外壁塗黑，盛水後放入一支木片 (12.5 cm x 2.5 cm)供雌蚊產卵，後來也有用條狀紙巾附貼於瓶子內壁來替代木條。這種誘蚊產卵

器廣為應用於世界各處埃及斑蚊的偵測與監測，甚至還將其作為防治病媒斑蚊之用。Fay and Eliason (1966) 比較誘蚊產卵器與幼蚊調查後認為二者可互補，且誘蚊產卵器更能經濟快速的偵測到埃及斑蚊產卵雌蚊的出現。Focks (2003)也認為此工具因容易布建、便宜、且相對敏感，確是調查埃及斑蚊族群的好工具。Chen (1973)於 1969 年將此工具用於新加坡國際機場埃及斑蚊的防治。進一步加上殺蟲網改良了誘蚊產卵器，後續 Lok, et al. (1977) 與 Cheng, et al. (1982)的研究也證實誘蚊產卵器確能用於實際防治登革熱的病媒蚊。甚至近年研究埃及斑蚊的產卵策略時，Chadee (2009)用的還是這種誘蚊產卵器。Richie, et al. (2004)所用的黏膠誘蚊器是由黑色塑膠筒，其內盛含 33%乾草浸液與 0.5 g 苜蓿丸的水 500 ml，筒內壁置兩片 15 x 5.5 cm 黏膠片所構成，當登革熱流行高蜂時，可捕獲高密度埃及斑蚊達 2~3.5♀/trap/week，施用殺幼蟲與成蟲藥劑防治後，所測密度下降為< 0.5♀/trap/week，他們認為此種黏性誘蚊器可作為監測產卵雌蚊的工具，也可用於評估防治成果與病毒的活動情形，而黏膠誘蚊器指數(Sticky ovitrap index=mean number of female per trap per week)更可用於登革熱傳播風險的量測估算，Williams, et al. (2006a, b) 就建議每戶設 4 個黏膠誘蚊器來作例行的監測。正如同 Focks (2003) 所強調的，精緻化現有的害蟲指標或發展更能反應疾病傳播風險的的新指標，其目的是要提供給社區、防治規劃甚至決策者更清楚與有意義的資訊。傳統的幼蚊調查以及使用的 HI、CI 或 BI 等指數，其實是顯示家戶孳生病媒斑蚊的數目或比率的。而各種誘引成蚊(尤其雌蚊)的工具與所獲指標則是用來估算有傳播登革熱風險的家戶數，也更適合用於孳生源調查無法發現幼蟲的時候。

本研究發現不同採集方法，雖然都捕獲較多之雌蚊，但僅

BG-stentinel 誘蚊器有顯著差異，誘蚊產卵器及背負式吸蟲機均捕獲相當數量之雄蚊（41.9%-42.5%），所以代表該地區亦為這些蚊蟲之孳生地，亦反應誘蚊產卵器對吸引雌蚊產卵並未有顯著效果。誘蚊產卵器捕獲之蚊蟲種類以熱帶家蚊為主，白線斑蚊次之，此次研究一個誘蚊器最高一週可誘到熱帶家蚊上百隻，雖然有些國外之研究亦有類似成果，但如何增進誘蚊產卵器吸引埃及斑蚊雌蚊來產卵，值得進一步研究。

本計畫未在冬天斑蚊中檢測出登革病毒，雖然檢測樣本數不多（埃及斑蚊 82♀44♂，白線斑蚊 2♀），但因我們使用的方法為靈敏度極高之 real-time RT-PCR 方法，其靈敏度可達 0.4684-7.8953 PFU/mL (Yang et al. 2010)，所以推測可能是因為南部地區冬天乾冷，不適合成蚊生存，成蚊於濕度低於 50%之死亡率高達 100%，所以帶病毒之雌蚊皆無法存活。另外冬天乾燥無雨水，大部份之卵無法孵化，靜待來年雨季再行孵化。此種結果再次呼應台灣地區登革熱非本土疾病，仍為旅遊傳染病（Shu et al. 2008, 2009, Chen et al. 2010, Kuan et al. 2010, Lin et al. 2010）。台灣地區之登革熱為每年自國外因旅遊而帶入病毒，再傳給當地病媒蚊，而引起每年的流行。

本計畫主要是以人口數做為誘蚊產卵器的放置數量，這是基於人口的多寡與登革熱傳播風險性的相關性而設計的，並且大規模在南台灣進行一年期的監測。相對於香港及新加坡則是依固定面積進行誘蚊產卵器的放置，但台灣在非都市區的人口分佈不平均，且地幅較為廣闊，利用人口作為調查依據可顯示其人與人媒介的重要性。總佈點位置為 5,180 個家戶，每週則有上萬筆資料需分析，人力物力的成本甚高，而資料的正確性與可靠性，更與志工、工作態度、居民配合度、回收資料處理流程與效率等因素息息相關。本計畫執行前的準備、各

項工作 SOP 擬定、教育訓練、溝通協調等均相當詳盡，但執行期間涉及多達 60 多位志工，仍有一些問題都在計畫執行過程中不斷地發生改進，不管是被動地志工在現場所面臨的問題，也主動的利用每週抽樣稽查方式發掘問題，讓計畫更加順利進行。

結論與建議

一、 本計畫設計的誘蚊產卵筒體積小，安置容易，且利用內置木棒或黏紙達到監測蚊卵、幼蟲或成蟲等不同目的。特別適用於登革熱好發地區，於低病媒密度或潛在孳生源不易發現時，誘蚊產卵器的使用，可替代孳生源稽查，有效偵測病媒蚊的發生。若配合衛星定位系統，則可進行疫區風險程度、疫病流行熱點或防治成效的評估。

二、 研究顯示本計畫使用的誘蚊產卵器如僅置放於室內(尤其於高度都市化地區)，則足以監測主要病媒埃及斑蚊的發生，並作為登革熱發生之預警。若以每 400 人放置一黏紙產卵筒的情形下，平均 100 個誘蚊產卵筒每週捕得病媒蚊 2.5 隻成蚊時，即應注意疫情暴發的可能性甚高，此法監測比現行衛生單位每月逢機部分區域的調查更為全面與客觀。

三、 本研究結果闡述闡述疫情中的兩種傳播模式—當地的環境因素傳播與人口移動的兩種不同的潛在傳播模式，在疫情的前、中、後期，處於相同的環境因素而感染的疫情傳播模式皆會存在，而因為人的移動造成感染的傳播模式，則主要會出現在疫情的後期。所以對於病例的居家週遭噴藥，僅能防治因當地的環境因素感染的傳播模式，對於人口移動造成的傳播模式並無改善的效果。最後，疫情務必於發生的早期及早阻斷，否則一旦人口移動以及相關複雜的因子後，將會使疫情的控制益發困難。

四、本計畫透過地理資訊系統(Geographic Information System, GIS)的協助，利用誘蚊產卵器所獲資料，得知在流行季時，高雄市三民區的東半部、高雄市苓雅區東半部以及高雄市前鎮區東北部，為病媒蚊分佈密度最高之處。高雄縣則發現鳳山市不論是流行季或非流行季，其病媒蚊穩定存在。另外林園鄉的南部，在流行季間病媒蚊的分佈亦屬密度較高的區域。

五、改善病媒蚊監測方法與技術—多年來台灣地方衛生單位對於登革熱病媒蚊每月都進行轄區部分地點的幼蚊調查，這種長期高成本的監測方式或有改善的空間。另一方面，不論病媒蚊族群季節性的消長如何，都以孳生源幼蚊的質性調查為主。事實上例如並無主要病媒埃及斑蚊發生的北部或東部地區，或可考量調整為二或三個月進行一次監測，又或許當布氏指數低於 3 或高於 30 後，可考量改用誘蚊產卵器來監測與掌握病媒的發生。

六、連結監測技術與防治策略—台灣每年自國外帶入登革病毒，傳給當地病媒蚊，而開使流行，因此疾病或病毒流行風險的探討其意義應不大，相對的掌握病媒孳生的時地變動，獲取正確的疫情資訊以及迅速的防治反應可能較為重要。更重要的思考或應是更精確的連結病媒密度與適當的防治策略，在不同的病媒發生量時施用不同的防治方法。

七、加強地方衛生單位彈性運用病媒監測技術的能力—因應各地區不同的都市化程度、人口密度、生活習慣、孳生容器類別、甚至財力物力等，應鼓勵並訓練其彈性運用適當監測的工具與技術。例如在病媒指數低但有病例發生時，或可使用黏膠成蚊器，一方面可監測傳病成蚊的發生與消長、另可鑑定病毒與直接捕殺前來產卵的雌成蚊，兼具防治功效。

八、建立病媒監測資料與流行風險系統—為即時反應病媒發生情形以及提供流行風險預警，宜建立透過網際網路，而地方與中央單位均可使用，甚至互動的平台，以利快速顯示病媒、疫情或流行風險的相關資訊。

九、若以誘蚊產卵器做為常態性的蚊蟲密度監測,如同香港及新加坡的方式，必需搭配現行孳生源的清除，才能有效監測蚊蟲實際密度，達到預先預警的功能。誘蚊器及吸蟲機則可在短時間內獲得蚊蟲密度。這些蚊蟲密度調查方法需要考慮施用的方法及時間，並對應於傳統指數，提出各種調查方法預警值的比較，以後可能利用這些方法來取代目前費時費工的調查方式。

十、利用各種病媒蚊指數與病例數分析,當月之病例數與當月病媒蚊密度或之後密度無關，但前一至三個月之布氏指數、住宅指數及背負式吸蟲機指數有正相關,所以在發生病例前一至三月就可以實施滅蚊計畫，預防登革熱發生或降低登革熱流行幅度。

十一、 誘蚊產卵器指數、背負式吸蟲機指數及傳統指數均呈現相同族群區趨勢，BG-Sentinel 誘蚊器配合乾冰放置對埃及斑蚊族群特別是低密度時，是一個較敏感的調查方法。

計畫重要研究成果及具體建議

一、 本計畫設計的誘蚊產卵器體積小，安置容易，且可利用內置木棒或黏紙達到監測蚊卵、幼蟲或成蟲等不同目的。若結合衛生定位系統對誘蚊產卵器置放地點定位，可進行疫區風險程度、疫病流行熱點或防治成效的評估。

二、 本計畫透過地理資訊系統(Geographic Information System, GIS)的協助，利用誘蚊產卵器調查數據據，得知在流行季時，

高雄市三民區的東半部、高雄市苓雅區東半部以及高雄市前鎮區東北部，為病媒蚊分佈密度最高之處，也是登革熱高風險地區。高雄縣鳳山市不論是流行季或非流行季，其病媒蚊密度穩定。林園鄉的南部，在流行季間病媒蚊的分布亦屬高風險區。

三、 提供高雄市及高雄縣埃及斑蚊的每年起始族群孳生地點 139 里，包括高雄市三民區 56 里、新興區 25 里、苓雅區 19 里、左營區 11 里、前鎮區 8 里、小港區 4 里、楠梓區 2 里、鹽埕區 1 里及高雄縣鳳山市 13 里，提供登革熱預防策略執行地點之參考。

四、 此計畫各種病媒蚊指數與病例數相關性分析，指出當月之病例數與前一至三個月之布氏指數、住宅指數及背負式吸蟲機指數有正相關，所以在發生病例前一至三月就可以實施滅蚊計畫，預防登革熱發生或降低登革熱流行幅度。

參考文獻

疾病管制局。97 年。登革熱防治工作指引。64 頁。

香港食物環境衛生署。登革熱。2007/12/23 摘自 http://www.fehd.gov.hk/safefood/dengue_fever/indexc.html。

Chen KL. The eradication of Aedes aegypti at the Singapore Paya Lebar International Airport. In Y. C. Chan, K. L. Chan and B. C. Ho (Eds.), *Vector control in southeast Asia*, 85-88. Ministry of Health, Singapore, 1973.

Cheng ML, Ho BC, Bartnett RE, Goodwin N. Role of a modified ovitrap in the control of *Aedes aegypti* in Houston, Texas, USA. *Bull. Wld Hlth Org.*, 60(2), 291-296, 1982.

Chadee DD. Ovipostion strategies adopted by gravid *Aedes aegypti* (L.) (Diptera: Culicidae) as detected by ovitraps in Trinidae, West Indies (2002-2006). *ACTA Tropica,* 111, 279-283, 2009.

Farrar J, Focks D, Gubler D, Barrera R, Guzman M G, Simmons C, Kalayanarooj S, Lum L, McCall P J, Lloyd L. Towards a global dengue research agenda. *Trop. Med. Int. Health*, 12, 695-699, 2007.

Fay, R. W. and Eliason, D. A. (1966). A preferred oviposition site as a surveillance method for *Aedes aegypti*. *Moq. News,* 26:531-535.

Fay RW, Pery AS. Laboratory studies of ovipositional preferences of *Aedes aegypti*. *Mosquito News*, 24, 276-281, 1965.

Focks DA. A review of entomological sampling methods and indicators for dengue vectors. *TDR/ IDE/Den/03.1,* Geneva, Switzerland: WHO, 40pp, 2003.

Kuan MM, Lin T, Chuang JH, Wu HS. Epidemiological trends and the effect of airport fever screening on prevention of domestic dengue fever outbreaks in Taiwan, 1998-27. Int. J. Infect. Dis.14: e693-7, 2010.

Lin CC, Huang YH, Shu PY, Wu HS, Lin YS, Yeh TM, Liu HS, Liu CC, Lei HY. Characteristic of Dengue Disease in Taiwan: 22–27Am. J. Trop. Med. Hyg. 82: 731-739, 2010.

Lok CK, Kiat NS, Koh TK. An autocidal ovitrap for the control and possible eradication of *Aedes aegypti*. *Southeast Asian J. Trop. Med. Publ. Health,* 8, 56-62, 1977.

Lloyd LS. Best practices for dengue prevention and control in the Americas. *Environmental Health Project, Strategic Report 7,* 106pp, 2003.

Mello AB, dos Santos WP, Rodrigues MAB, Candeias ALB, Gusmão CMG (2008) Image Segmentation of Ovitraps for Automatic Counting of *Aedes aegypti* Eggs. 30th Annual International IEEE EMBS Conference. Vancouver, British Columbia, Canada, August 20-24, 2008

Morrison AC, Zielinski-Gutierrez E, Scott TW, Rosenberg R (2008) Defining challenges and proposing solutions for control of the virus vector *Aedes aegypti*. PLoS Med 5: e68.

Ooi EE, Goh KT, Gubler DJ (2006) Dengue prevention and 35 years of vector control in Singapore. Emerging Infectious Diseases 12(6): 887- 883.

Regis L, Monteiro AM, de Melo-Santos MAV, Silveira Jr JC, Furtado AF, Acioli RV, Santos GM, Nakazawa M, Carvalho MS, Ribeiro Jr PJ, de Souza WV. Developing new approaches for detecting and preventing *Aedes aegypti* population outbreaks: basis for surveillance,alert and control system. Mem Inst Oswaldo Cruz, Rio de Janeiro 103: 50-59, 2008.

Richie SA, Long S, Smith G, Pyke A, Knox TB. Entomological investigations in a focus of dengue transmission in Cairns, Queensland, Australia, by using the sticky ovitraps. *J. Med. Entomol.*, 41(1), 1-4,

2004.

Shu PY, Su CL, Liao TL, Yang CF, Chang SF, Lin CC, Chang MC, Hu HC, Huang JH. Molecular Characterization of Dengue Viruses Imported Into Taiwan during 23–27: Geographic Distribution and Genotype Shift Am. J. Trop. Med. Hyg., 8:139-146, 2009.Kuan , 2010.

Taiwan CDC. Title. Notifiable Infectious Diseases Statistics System(http://nidss.cdc.gov.tw/SingleDisease.aspx?Pt=s&Dc=1&Dt=2&disease=61), Down load at 11/14/2010.

WHO. Dengue and dengue haemorrhagic fever. (http://www.who.int/mediacentre/factsheets/fs117/en). Down load at 11/14/2010.

Yang CF, Shu PY, Chen CF, Su CL, Teng HJ, Lu LC, Lin C, Wang CY, Huang JH, Wu HS. 2010. Screening of Mosquito Infections Using SYBR Green I-based Real time RT-PCR with Group Primers of Flavivirus and Alphavirus in Taiwan. Journal of Virological Methods 168:147-151, 2010.

Williams CR, Long SA, Russell RC, Ritchie SA. Field efficacy of the BG-Sentinel compared with CDC Backpack Aspirators and CO2-baited EVS traps for collection of adult Aedes aegypti in Cairns, Queensland, Australia. J. Am. Mosq. Control Assoc. 22:296–300, 2006a.

Williams CR, Long SA, Russell RC, Ritchie SA. Optimizing ovitrap use for *Aedes aegypti* in Cairns, Queensland, Australia: Effects of some abiotic factors on field eefficacy. J. Am. Mosq. Control Assoc. 22:635–640, 2006b.

研究區域偵測點分佈情形



圖一、本計畫病媒蚊監測點設置的空間分佈。



(B)



(C)



圖二、高雄氣候（A）、誘蚊產卵器雌蚊每週變動圖（B）及傳統指數（布氏指數、住宅指數及成蚊指數）每週變動圖（C）。



圖三、98年6月14日至99年6月13日（53週）誘蚊產卵器產卵棒戶內及戶外陽性率週消長趨勢圖（A）高雄縣市（B）高雄市及（C）高雄縣。



圖四、98 年 6 月 14 日至 99 年 6 月 13 日（53 週）以誘蚊產卵器產卵棒所採集之卵孵化後斑蚊總數（A）與高雄市（B）及高雄縣（C）埃及斑蚊及白線斑蚊幼蟲週消長趨勢圖。



圖五、98 年 6 月 14 日至 99 年 6 月 13 日（53 週）高雄市誘蚊產卵器塑膠黏片所採集之斑蚊數（A）、埃及斑蚊百分比（B）及每百個誘蚊產卵器採集之斑蚊數（C）。



圖六、98 年 6 月 14 日至 99 年 6 月 13 日（53 週）高雄縣誘蚊產卵器塑膠黏片所採集之斑蚊數（A）、埃及斑蚊百分比（B）及每百個誘蚊產卵器採集之斑蚊數（C）。



圖七、98 年 6 月至 99 年 3 月 7 日高雄市 11 個行政區及高雄縣 6 個鄉鎮市區(鳳山市、梓官鄉、林園鄉、仁武鄉、大寮鄉及鳥松鄉)誘蚊產卵器監測之埃及斑蚊陽性次數圖及密度圖。



圖八、99 年 1 月至 3 月 7 日高雄市 11 個行政區及高雄縣 6 個鄉鎮市區(鳳山市、梓官鄉、林園鄉、仁武鄉、大寮鄉及鳥松鄉)誘蚊產卵器監測之埃及斑蚊陽性次數圖及密度圖。



圖九、2009 年 7 月起至 2010 年 1 月止，本土病例發病週次分布。

研究區域本土病例分佈



圖十、 2009 年 7 月至 2010 年 1 月止的本土病例空間分佈



圖十一、每一個病例與其他病例發病相隔日數以及間隔距離。X 軸為間隔距離，單位為 100 公尺。Y 軸為發病相隔日數，單位為天。顏色越接近暖色系，代表該座標位置上存在病例-病例之間的配對越多。





圖十二、四個時空聚集區，與其他病例關聯最密集的前 10%病例。







圖十三、第一時空聚集區與第四時空聚集區病例發病週次分佈(括號為研究週次)



圖十四、第一與第四時空聚集病例，病例發生前四週且 1 公里內偵測點涵蓋平均卵數。



圖十五、2009 年高雄地區登革熱病例發生月份消長圖（A）及各季間病媒蚊所誘集卵數（B）。

（A）



（B）



（C）



圖十六、2009 年高雄地區登革熱病媒蚊空間分布圖（A）及病媒蚊群聚（B）。

（A）



（B）



圖十七、2009 年高雄地區登革熱風險區估算（2009 年登革熱風險區=病媒蚊流行高峰 11 月份空間分布圖 x 人口密度）（A）及 2009 年預估風險群聚區塊、2009 年 8 月至 2010 年 1 月高雄地區登革熱發生空間分布圖及發生群聚區塊（A）。





圖十八、2009 年 8 月至 2010 年 1 月高雄地區登革熱發生空間分布圖（A）及發生群聚區塊（B）。



圖十九、高雄市前鎮區及高雄縣鳳山市為各種採樣方法主要的調查區域。左圖為此兩區域之所在位置，右圖分散之小黑點為誘蚊產卵器放置位置。

(A)



(B)



圖二十、各種取樣方法指數之每月趨勢圖(A)及氣候變化趨勢(B)。(A)誘蚊器指數(BG trap index)、吸蟲機指數(Backpack aspirator index)及本土登革熱病例數(Local dengue cases)皆除以 10 呈現。左邊 Y 軸以百分比(%)為單位，右邊 Y 軸以數字為單位(包含布氏指數 Breateu index 及本土登革熱病例數)。(B) 每月平均溫度(Temperature)及雨量(Rainfall)。

表一、研究地點高雄市及部份高雄縣各鄉鎮市區面積、人口密度、登革熱病媒蚊分布百分比及誘蚊產卵器放置家戶數。

縣市	鄉鎮市區	面積（km^2）	97 年密度（人/km^2）	埃及斑蚊%	白線斑蚊%	分層區塊	誘蚊產*卵器放置家戶
高雄市	新興區	1.98	28,691	94.1%	5.9%	1	125
高雄市	苓雅區	8.15	22,910	95.4%	4.6%	1	431
高雄市	旗津區	1.46	20,452	92.1%	7.9%	1	69
高雄市	鹽埕區	1.42	19,820	94.9%	5.1%	1	61
高雄市	三民區	19.79	17,988	94.6%	5.4%	1	849
高雄市	前金區	1.86	16,031	94.2%	5.8%	1	64
高雄市	前鎮區	19.13	10,486	90.1%	9.9%	1	475
高雄市	楠梓區	25.83	6,598	82.9%	17.1%	1	405
高雄市	左營區	19.39	9,731	66.6%	33.4%	2	454
高雄市	鼓山區	14.75	8,520	73.9%	26.1%	2	297
高雄市	小港區	39.86	3,842	67.0%	33.0%	2	365
高雄縣	鳳山市	26.75	12,684	67.1%	32.9%	2	807
高雄縣	梓官鄉	11.60	3,188	27.8%	72.2%	2	86
高雄縣	仁武鄉	26.59	2,595	30.4%	69.6%	2	166
高雄縣	林園鄉	32.29	2,198	32.8%	67.2%	2	164
高雄縣	鳥松鄉	24.59	1,697	25.2%	74.8%	2	100
高雄縣	大寮鄉	71.04	1,541	27.5%	72.5%	2	262
總計		346.48	---	---	---	---	5,180

*每家戶放置 2 個誘蚊產卵器，戶內客廳放置 1 個，戶外放置 1 個。

表二、塑膠黏片與產卵棒實驗室實驗結果。

項目	重複數	5對埃及斑蚊					10對埃及斑蚊				
		誘捕成蚊數	雌蚊黏捕率	雄蚊黏捕率	斑蚊黏捕率	卵數	誘捕成蚊數	雌蚊黏捕率	雄蚊黏捕率	斑蚊黏捕率	卵數
塑膠黏片與產卵棒	I	3♀2♂	60%	40%	50%	1	5♀5♂	50%	50%	50%	0
	II	2♀3♂	40%	60%	50%	30	6♀5♂	60%	50%	55%	38
產卵棒	I	–	–	–	–	59	–	–	–	–	326
	II	–	–	–	–	243	–	–	–	–	345

表三、98年6月至7月世運22個場點誘蚊產卵器產卵棒及黏紙前試驗結果。

週別	誘蚊器數目	產卵棒				塑膠黏片			
		戶內陽性率	戶外陽性率	平均陽性率	總卵數	戶內陽性率	戶外陽性率	平均陽性率	總斑蚊數
第一週	44	0.0	36.4	18.2	237	4.5	13.6	9.1	4
第二週	44	0.0	13.6	6.8	109	0.0	31.2	15.6	10
第三週	44	0.0	18.2	9.1	268	0.0	13.6	6.8	7
第四週	44	0.0	13.6	6.8	171	4.5	22.7	13.6	7
第五週	44	0.0	27.3	13.7	200	4.5	22.7	13.6	8

表四、98 年 6 月至 99 年 6 月誘蚊產卵器誘集之卵數與斑蚊成蚊數之處理間比較。

處理	n	每個誘蚊產器捕獲卵數 平均±SD	誘蚊產器採集卵陽性率 平均 ± SD	每個誘蚊產器捕獲斑蚊數 平均± SD	誘蚊產器捕獲斑蚊陽性率 平均±SD
產卵棒	204	0.69a±0.65	2.63a ±2.24	-----	
產卵棒加塑膠黏片	204	0.08b±0.09	0.50b ±0.48	2.95 a±1.36	2.28a±1.27
塑膠黏片	204	-----	-----	1.90 b±2.50	1.63b±1.71

*Means within each column having the same letter are not significantly different at P≥0.05.

表五、99 年 1 月至 3 月 7 日冬天埃及斑蚊分布重點區塊。

縣市別	市區別	里別
高雄市	三民區	安泰里、安邦里、民享里、十美里、德行里、德東里、鼎勝里、鼎強里、鼎力里、鼎金里、鼎泰里、鼎中里、鼎西里、灣愛里、灣子里、灣興里、灣復里、灣成里、灣華里、灣勝里、灣利里、灣中里、本和里、本館里、本元里、本武里、本文里、本安里、本上里、本揚里、寶龍里、寶珠里、寶興里、寶中里、寶玉里、寶泰里、寶安里、寶盛里、寶德里、寶獅里、寶國里、寶慶里、寶業里、寶華里、寶民里、安生里、安吉里、安寧里、安康里、安發里、博惠里、港新里、同德里、德仁里、港北里、港東里
	新興區	順昌里、文昌里、開平里、興昌里、光耀里、德生里、南港里、中東里、成功里、德政里、德望里、華聲里、永寧里、新江里、黎明里、愛平里、東坡里、漢民里、玉衡里、秋山里、明莊里、大明里、榮治里、仁聲里、蕉園里
	苓雅區	正道里、正心里、正大里、正文里、正言里、永康里、五福里、民主里、五權里、福南里、奏捷里、安祥里、凱旋里、尚義里、中正里、林安里、城西里、意誠里、衛武里
	左營區	菜公里、新光里、新下里、埤東里、華豐里、新上里、果貿里、果惠里、果峰里、裕豐里、裕興里
	前鎮區	瑞華里、瑞昌里、瑞興里、瑞崗里、瑞隆里、瑞東里、瑞和里、瑞平里
	小港區	大苓里、正苓里、二苓里、三苓里
	楠梓區	國昌里、裕昌里
	鹽埕區	博愛里
高雄縣	鳳山市	善美里、一甲里、富甲里、鎮南里、埤頂里、美齡里、忠孝里、忠義里、中和里、協和里、新強里、興中里、鎮西里

表六、98 年 6 月至 99 年 6 月高雄市前鎮區及高雄縣鳳山市捕獲成蚊之蚊種組成 [隻數(總數百分比)]

蚊蟲種類	誘蚊產卵器			BG-Sentinel 誘蚊器			背負式吸蟲機		
	總數	雌蚊數	雄蚊數	總數	雌蚊數	雄蚊數	總數	雌蚊數	雄蚊數
埃及斑蚊	1,136 (5.7%)	679 (5.9%)	457 (5.4%)	265 (45.8%)	170 (40.6%)	95 (59.4%)	342 (26.7%)	173 (23.3%)	169 (31.5%)
白線斑蚊	227 (1.1%)	189 (1.7%)	38 (0.4%)	14 (2.4%)	10 (2.4%)	4 (2.5%)	34 (2.7%)	26 (3.5%)	8 (1.5%)
無法辨別斑蚊	517 (2.6%)	298 (2.6%)	219 (2.6%)	0 (0.0%)	0 (0%)	0 (0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
斑蚊總數	1,880 (9.5%)	1,166 (10.2%)	714 (8.5%)	279 (48.2)	180 (43.0%)	99 (61.9%)	376 (29.4%)	199 (26.8%)	177 (33.0%)
白腹叢蚊	40 (0.2%)	22 (0.2%)	18 (0.2%)	93 (16.1)	84 (20.0%)	9 (5.6%)	130 (10.2%)	87 (11.7%)	43 (8.0%)
黃尾家蚊	1 (0.009%)	1 (0.009%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
熱帶家蚊	17,973 (90.3%)	10,246 (89.6%)	7,727 (91.3%)	207 (35.8)	155 (37.0%)	52 (32.5%)	773 (60.4%)	457 (61.5%)	316 (59.0%)
總　計	19,894	11,435	8,459	579	419	160	1279	743	536

表七、98 年 6 月至 99 年 6 月高雄市前鎮區及高雄縣鳳山市各種採樣指數比較表。

調查月份	布氏指數	住宅指數	成蚊指數 (x100)	誘蚊產卵器指數	BG sentinel 誘蚊器指數	背負式吸蟲機指數	登革熱病例數
98.06	5.2	4.1%	1.0	3.2%	50.0%	30.0%	0
98.07	6.9	5.4%	1.8	5.3%	100.0%	60.0%	1
98.08	7.7	5.4%	1.5	4.6%	62.5%	55.0%	5
98.09	8.4	6.0%	2.2	6.1%	62.5%	67.5%	20
98.10	7.0	5.2%	2.6	5.5%	50.0%	55.0%	88
98.11	6.0	4.6%	2.5	3.5%	50.0%	37.5%	154
98.12	2.7	2.2%	1.0	2.2%	50.0%	12.5%	99
99.01	1.1	1.0%	0.5	0.5%	25.0%	8.3%	7
99.02	0.9	0.7%	0.5	0.7%	37.5%	1.7%	0
99.03	0.9	0.8%	0.5	1.2%	0.0%	0.0%	0
99.04	1.0	0.9%	0.7	1.1%	0.0%	13.3%	0
99.05	1.9	1.6%	0.8	1.4%	75.0%	13.3%	0
99.06	6.1	4.8%	3.2	2.1%	100.0%	23.3%	0
平均	4.3	3.3%	2.7	2.9%	51.0%	29.0%	---

備註：誘蚊產卵器指數、BG-Sentinel誘蚊器指數及吸蟲機指數為捕獲斑蚊陽性戶數的比例(等於傳統指數中之住宅指數或住宅陽性率)。

表八、各種採集方法之相關性(Pearson Correlation)分析。

變數	住宅指數	成蚊指數	誘蚊產卵器指數	BG-Sentinel誘蚊器指數	背負式吸蟲機指數
布氏指數	**r=0.997** **p=0.00**	r=0.42 p=0.15	**r=0.93** **p=0.00**	**r=0.65** **p=0.02**	**r=0.94** **p=0.00**
住宅指數	---	r=0.46 p=0.11	**r=0.92** **p=0.00**	**r=0.68** **p=0.01**	**r=0.93** **p=0.00**
成蚊指數	---	---	r=0.48 p=0.09	**r=0.58** **p=0.04**	r=0.51 p=0.07
誘蚊產卵器指數	---	---	---	r=0.50 p=0.08	**r=0.98** **p=0.00**
BG-Sentinel誘蚊器指數	---	---	---	---	r=0.54 p=0.057

備註：誘蚊產卵器指數、BG-Sentinel誘蚊器指數及吸蟲機指數為捕獲斑蚊陽性戶數的比例(等於傳統指數中之住宅指數或住宅陽性率)。

表九、各種採集方法與病例數之相關性(Pearson Correlation)分析。

	當月登革熱病例數	前1月病例數	前2月病例數	前3月病例數	前4月病例數
布氏指數	r=0.24	**r=0.57**	**r=0.71**	**r=0.67**	r=0.47
	p=0.43	**p=0.04**	**p=0.01**	**p=0.01**	p=0.11
住宅指數	r=0.25	**r=0.56**	**r=0.67**	**r=0.64**	r=0.49
	p=0.41	**p=0.046**	**p=0.01**	**p=0.02**	p=0.09
成蚊指數	r=-0.07	r=-0.01	r=0.04	r=0.38	**r=0.72**
	p=0.82	p=0.98	p=0.90	p=0.21	**p=0.006**
誘蚊產卵器指數	r=0.26	**r=0.66**	**r=0.81**	**r=0.69**	r=0.49
	p=0.38	**p=0.01**	**p=0.001**	**p=0.01**	p=0.09
BG-Sentinel 誘蚊器指數	r=-0.01	r=0.07	r=0.19	r=0.38	r=0.45
	p=0.98	p=0.83	p=0.53	p=0.21	p=0.12
背負式吸蟲機指數	r=0.19	**r=0.61**	**r=0.80**	**r=0.74**	r=0.53
	p=0.53	**p=0.03**	**p=0.001**	**p=0.004**	p=0.07
當月登革熱病例數	---	**r=0.67**	r=0.08	r=-0.23	r=-0.29
		p=0.01	p=0.79	p=0.45	p=0.33

表十、誘蚊產卵器放置地點稽查時間與成果。

稽查時間	稽查單位	稽查地點	稽查戶數	室外合格率(%)	室內合格率(%)
98/07/02	屏科大	林園鄉	9	66.7	71.4
98/07/03	屏科大	三民二區	12	83.3	91.7
98/07/09	屏科大	鳳山市	30	7.1	17.3
98/07/10	屏科大	鼓山區	10	100.0	100.0
98/07/15	屏科大	三民一區	32	81.3	78.1
98/08/20	屏科大	鳳山市	22	16.7	17.7
98/08/27	屏科大	鳳山市	27	39.1	43.5
98/08/28	屏科大	左營區	10	90.0	80.0
98/08/28	屏科大	鳳山市	18	25.0	25.0
98/09/03	屏科大	鳳山市	25	72.0	72.0
98/09/04	屏科大	楠梓區	11	81.8	54.6
98/09/04	屏科大	鳳山市	27	68.9	68.9
98/09/09	屏科大	小港區	31	79.2	79.2
98/09/11	屏科大	三民二區	5	100.0	100.0
98/09/16	屏科大	小港區	24	95.0	95.0
98/09/28	屏科大	前鎮區	4	75.0	50.0
98/10/02	屏科大	苓雅區	24	50.0	41.7
98/10/07	屏科大	大寮鄉	23	80.0	80.0
98/10/14	屏科大	大寮鄉	18	100.0	100.0
98/10/22	屏科大	梓官鄉	4	100.0	100.0
98/10/23	屏科大	旗津區	29	90.0	90.0
98/10/29	屏科大	旗津區	46	70.4	70.4
98/11/05	屏科大	前金區	7	100.0	28.6
98/11/05	屏科大/疾管局	鳳山市	10	66.7	62.5
98/11/05-06	屏科大/疾管局	前鎮區	10	77.8	77.8
98/11/06	屏科大/疾管局	旗津區	10	100%	90%
99/04/15	屏科大/疾管局	鼓山區	10	90.0	40.0
99/04/15	屏科大/疾管局	三民區	10	60.0	10.0
99/04/16	屏科大/疾管局	鳳山市	10	30.0	30.0
99/07/26	屏科大/疾管局	苓雅區(回收)	10	90.0	100.0
99/07/27	屏科大/疾管局	前金區(回收)	10	100.0	100.0
99/07/27	屏科大/疾管局	鳳山市(回收)	10	90.0	90.0

表十一、誘蚊產卵器放置成本效益評估。

項目	單價	放置53週	放置10,360個誘蚊產卵器53週
誘蚊產卵器監測			
誘蚊產卵器	54.20	54.20	561,512
塑膠黏片	2.63	139.39	1,444,080
產卵棒	0.40	21.20	219,632
殺幼蚊劑	2.04	108.12	1,120,123
放置及回收	20.00	1,060.00	10,981,600
合計	79.27	1,382.91	14,326,947

緊急防治費用

(1 組工作 4 小時，噴 20 戶 4-5 層獨棟，警察、噴霧機、鎖匠等費用)

項目	數量	單價	1 個病例 (噴半徑 50 公尺，約 40-80 戶，需 2-4 組)
前導人員	1 人	800	1,600-3,200
孳生源清除人員	1 人	800	1,600-3,200
噴藥領隊	1 人	800	1,600-3,200
噴藥工	2 人	1,600	3,200-6,400
殺蟲劑	1 瓶	2,000	4,000-8,000
合計	---	6,000	12,000-24,000

附錄一


2010 The Internation Forum for
Dengue Control
登革熱防治國際論壇
Kaohsiung, Taiwan Nov. 25~27, 2010
時間
地點
2010/11.25~26
高雄市蓮潭國際文教會館
高雄市左營區崇德路801號 TEL:07-2354999
主辦單位 行政院衛生署疾病管制局
承辦單位 屏東科技大學
協辦單位 台灣大學、元培科技大學
中興大學、嘉南藥理科技大學
高雄大學、大仁科技大學
屏東科技大學植物醫學系
連絡電話：08-7703202 轉 6168
連絡地址：912屏東縣內埔鄉學府路一號

2010 登革熱防治國際論壇

時間：中華民國 99 年 11 月 25~26 日

地點：高雄市蓮潭國際文教會館（高雄市左營區崇德路 801 號，TEL:07-2354999）

議程：

11 月 25 日 (THURSDAY, Nov 25, 2010)

Time	Activity	Speaker
08:30~09:30	報到 Registration and Information	
09:30~09:45	開幕 Opening remark 長官致詞	行政院衛生署疾病管制局 CDC official
09:45~10:00	團體照 Taking Group photo	
第一節：亞洲地區登革熱之防治（主持人:徐爾烈教授） Section 1: The control of dengue fever in Asia (Moderator : Dr. Err-Lieh Hsu)		
10:00~10:30	臺灣的流行病學探究如何有助於全球登革出血熱疫情的控制 How could global control of dengue hemorrahgic fever achieve success - Taiwan's valuable experiences from epidemiological findings	Dr. Chwan-Chuen King 金傳春 Institute of Epidemiology, College ofPublic Health, National Taiwan University. Taipei, Taiwan
10:30~11:00	茶敘時間 Break	
11:30~12:00	越南的登革熱：現況、策略方向與挑戰 Dengue in Vietnam: Current situation - Strategic direction - Challenge	Dr. Tran Vu Phong Representative of Medical Entomology and Zoology National Institute of Hygiene and Epidemiology, Vietnam
11:30~12:00	廣東省登革熱流行與控制現狀及面臨挑戰 Epidemics and Control of Dengue Fever in Guangdong Province and its Challenges	Dr. L. F. Lin Guangdong Center for Disease Control and Prevention, China
12:00~13:30	午餐 Lunch Break	
第二節：亞洲地區登革熱病媒之防治（主持人:金傳春教授） Section 2: The control of dengue fever in Asia (Moderator : Dr. Chwan-Chuen King)		
13:30~14:00	以菲律賓馬尼拉公立醫院斑蚊監測作為登革熱病媒防治的基礎 *Aedes* surveillance of selected public hospitals in Metro Manila Philippines as basis for developing vector control.	Dr. Estrella Irlandez Cruz DAPE, MPH Department of Medical Entomology Research Institute for Tropical Medicine, Philippines

14:00~14:30	蘇力菌 (AM 65-52)對登革熱病媒蚊之防治 *Bacillus thuringiensis israelensis* (strain AM 65-52) for the control of dengue vectors.	Dr. Seleena Benjamin Valent BioSciences Corp Malaysia Dr. Hsiu-hua Pai 白秀華 College of Humanities and Social Sciences, National university of Kaohsiung, Taiwan
14:30~15:00	病原真菌用於斑蚊防治之潛力 The potential of using entomopathogenic fungi in *Aedes* control	Dr. Li-Cheng Tang 唐立正 Department of Entomology, National Chung-Hsing University, Taichung, Taiwan
15:00~15:30	茶敘時間 Tea Time	
第三節： 台灣地區登革熱病媒化學防治及其抗藥性之發生（主持人:張念台教授） Section 3: Chemical control and insecticides resistance of dengue vectors in Taiwan (Moderator : Dr. Niann Tai Chang)		
15:30~16:00	台灣登革熱防治與埃及斑蚊之抗藥性 Dengue fever control and insecticides resistance in *Aedes aegypti* in Taiwan.	Dr. Err-Lieh Hsu 徐爾烈 Dept. Entomology, National Taiwan Univ.Taipei, Taiwan Dr. Huai Hui ,Wu 吳懷慧 Department of Biotechnology, Tajen University, Pingtung, Taiwan Dr. Yi-Pey Luo 羅怡珮 Department of Biotechnology, Chia-Nan University of Pharmacy and Science, Tainan, Taiwan
16:00~16:30	以鈉通道基因突變頻率預測埃及斑蚊對除蟲菊脂類藥劑的抗性 Prediction of pyrethroid resistance by frequency of sodium channel gene mutations in *Aedes aegypti*	Dr. Shu-Mei Dai 戴淑美 Department of Entomology, National Chung-Hsing University, Taichung, Taiwan Dr. Ying-Hsi Lin 林鶯熹 Department of Medical Laboratory Science and Biotechnology, Yuanpei University, HsinChu, Taiwan
16:30~17:00	台灣南部地區登革熱及病媒蚊防治整合計畫報告 Report on the integrated research program for the management of dengue epidemics and vector mosquitoes in Southern Taiwan	Dr. Niann Tai Chang 張念台 Department of Plant Medicine, Pingtung University of Science and Technology, Taiwan

FRIDAY, Nov 26, 2010

Time	Activity	Speaker
08:30~09:00	報到 Registration and Information	
第四節:登革熱流行之監測及預測（主持人:鄧華真科長） Section 4: Surveillance and prediction of dengue epidemics (Moderator : Dr. Hwa-Jen Teng)		
09:00~09:30	台灣登革熱高危險區登革熱病媒蚊採樣方法評估 Evaluation of collection methods of dengue vectors in dengue high-risk areas of Taiwan	Dr. Hwa-Jen Teng 鄧華真 Chih-Yuan Wang 王智源 Research and Diagnostic Center, Centers for Disease Control, Taipei, Taiwan
09:30~10:00	建置以地理資訊為基礎的登革熱病媒蚊監測	Dr. Tzai-Hung Wen, 溫在弘

	系統於人蚊互動之時空分析 Establishment of a GIS-based Mosquito Surveillance System for Modeling Human-Vector Interactions in Time and Space	Department of Geography, National Taiwan University, Taipei, Taiwan. Dr. N. T. Chang 張念台 Pingtung University of Science and Technology, Taiwan Dr. Hwa-Jen Teng 鄧華真 Research and Diagnostic Center, Centers for Disease Control, Taipei, Taiwan
10:00~10:30	境外移入病例在本土登革流行初期的起始作用：1998-2007 的時序分析 The Role of Imported Cases in Initiation Dengue Epidemics in Taiwan: Temporal Analyses of Epidemiological and Meteorological Factors of Dengue during 1998-2007	Dr. Chuin-Shee Shang 尚君璽 Division of Environmental Health and Occupational Medicine, National Health Research Institutes, Taipei, Taiwan Dr. Chi-Tai Fang 方啟泰, Dr. Chung-Ming Liu 柳中明, Dr. Tzai-Hung Wen 溫在弘, Dr. Kun-Hsien Tsai 蔡坤憲, Dr. Chwan-Chuen King 金傳春 National Taiwan University, Taipei, Taiwan,
10:30~11:00	茶敘時間 Break	
第五節:登革熱病毒與病媒（主持人:杜武俊副教授） Section 5: Dengue viurs and vectors (Moderator : Dr. Wu-Chun Tu)		
11:00~11:30	蚊子對登革熱病毒之媒介 Vector competence of mosquitoes for dengue virus	Dr. Wu-Chun Tu 杜武俊 Department of Entomology, National Chung-Hsing University, Taichung, Taiwan
11:30~12:00	以遺傳策略防治登革熱病毒 Establishing a gene driver for preventing Dengue virus	Dr. C. H. Chen 陳俊宏 National Health Research Institutes, Taipei, Taiwan
12:00~13:30	午餐 Lunch Break	
第六節:台灣登革熱及病媒蚊防治實務（主持人:CDC 長官） Section 6: Practice of Dengue and Dengue vectors Control in Taiwan (Moderator : CDC official)		
13:30~14:00	高雄市登革熱防治現況 Report and Discussion of Dengue Control in Kaohsiung City	1.衛生署疾管局第五分局 The Fifth Branch Office of CDC, Department of Health, Kaohsiung 2.高雄市政府衛生局疾病管制處 CDC of Kaohsiung City Health Department
14:00~14:30	高雄縣登革熱防治現況 Report and Discussion of Dengue Control in Kaohsiung County	高雄縣政府衛生局 The Health Bureau of Kaohsiung County Government
14:30~15:00	茶敘時間 Tea Time	

15:00~15:30	屏東縣登革熱防治現況 Report and Discussion of Dengue Control in Pingtung County	屏東縣政府衛生局 The Health Bureau of Pingtung County Government
15:30~16:00	台南縣市登革熱防治現況 Report and Discussion of Dengue Control in Tainan City and Tainan county	台南縣市政府衛生局 CDC of Tainan City (Tainan county) Health Department
16:00~16:30	討論 panel discussion	
16:30	閉幕 Closing	

台灣登革熱高危險區登革熱病媒蚊採樣方法評估

行政院衛生署疾病管制局研究檢驗中心 鄧華真、王智源

埃及斑蚊與白線斑蚊是台灣地區的登革熱主要病媒蚊。由各種採樣方法所產生代表當地斑蚊族群指標為評估疾病發生風險及防治措施的決定性因素。我們在高雄縣鳳山市及高雄市前鎮區進行一年的調查，比較傳統斑蚊指數(布氏指數、住宅指數及成蚊指數)、改良之誘蚊產卵器、BG-Sentinel™誘蚊器及 CDC 背負式吸蟲機器等五種採樣方法。這些指數（除 BG-Sentinel™誘蚊器外）之月消長均相似，同溫度及降雨量之變化，六月增加，至七-九月高峰，十月下降，一月至五月谷底。BG-Sentinel™誘蚊器除了三月及四月沒有誘集到斑蚊，其他月份均捕獲相當數量。布氏指數、住宅指數、誘蚊產卵器指數及背負式吸蟲機指數間有很高的顯著性正相關性（$r>0.91$, $p<0.000$）。誘蚊器與傳統斑蚊指數間亦有顯著性正相關（$r=0.581$-0.6793, $p<0.05$）。另外，當地本土病例數與布氏指數、住宅指數及背負式吸蟲機有前移一至三個月之正相關($r=0.6668$-0.8053, $p<0.05$)。我們的結論為發生病例前一至三月就要實施滅蚊計畫及 BG-Sentinel™誘蚊器是一個敏感的調查方法，特別是低密度時。

Evaluation of collection methods of dengue vectors in dengue high-risk areas of Taiwan

Hwa-Jen Teng ,Chih-Yuan Wang, Research and Diagnostic Center, Centers for Disease Control, Taiwan, ROC

Abstract: *Aedes* aegypti and *Ae. albopictus* are the main vectors of dengue in Taiwan. The indicators generated by various sampling methods to estimate the local population of *Aedes* mosquitoes are critical for evaluating the disease-risk and the effectiveness of control measures. Five sampling methods, traditional *Stegomyia* indece (Breteaux index, house index and adult index), modified ovitraps, BG-Sentinel™ mosquito traps and CDC backpack aspirators, were compared for one year in Fensheng City, Kaohsiung County and Chichen District, Kaohsiung City. Monthly fluctuation

of all indice (except for BG-Sentinel trap index) presented similar pattern, which were concurred with temperatures and rainfalls of the same year. BG-Sentinel traps collected some number of *Aedes* mosquitoes except for March and April. Breteau index, house index, ovitrap index and backpack aspirator index had significantly higher positive correlation with each other ($r>0.91$, $p<0.000$). BG-Sentinel trap index had significantly positive correlations with *Stegomyia* indice ($r=0.581$-0.6793, $p<0.05$). Additionally, the number of indigenous dengue cases had 2 month time-forward effect with Breteau index, house index, and backpack aspirator index ($r=0.6668$-0.8053, $p<0.05$). In conclusion, mosquito control measures should be conducted 1-3 months before the cases was found and BG-Sentinel™ trap was a sensitive collection method, even in a low population.

建置以地理資訊為基礎的登革熱病媒蚊監測系統於人蚊互動之時空分析

溫在弘[1]、張念台[2]、鄧華真[3]

1. 國立台灣大學地理環境資源學系
2. 國立屏東科技大學植物醫學系
3. 衛生署疾病管制局研究檢驗中心

為了更清楚瞭解登革熱病媒蚊在登革熱高發生率地區的族群動態，本研究針對高雄市全市及高雄縣七個行政區（鳳山市、梓官鄉、仁武鄉、鳥松鄉、大寮鄉、林園鄉），依照該地區人口密度特性與病媒蚊棲息環境來部署病媒蚊偵測點。本研究區總共設置 5180 個偵測點，每週持續收集病媒蚊資料，包括：蟲卵數、埃及斑蚊與白線斑蚊幼蟲數等數據。資料收集期間為 2009 年 6 月 7 日起至 2010 年 6 月 13 日。透過一年內逐週的病媒蚊資料收集，輔以網路地理資訊平台 Google Maps，本研究建立「病媒蚊監測地圖系統」(**GIS-based Mosquito Surveillance Mapping System**)，將提供衛生單位掌握病媒蚊連續監測的時間與空間分佈趨勢，透過連續監測資料呈現病媒蚊活動頻繁的區域，即為登革熱病媒防治重點之所在。此外，由於感染登革熱病例，亦是造成病毒傳播過程的重要因子，因此本研究亦分析從 2009 年 7 月起至 2010 年 1 月止，登革熱本土病例在時間與空間聚集的情形，透過了解病例時間與空間的聚集情形，以期能釐清整體疫情的脈絡，進而掌握疫情發展與擴散的模式而有最佳的介入措施。本研究建立病媒蚊監測資料與病例之間的時間與空間關聯性，有助於描繪人蚊接觸與互動對於整體疫情發展的影響。

Establishment of a GIS-based Mosquito Surveillance Mapping System for Modeling Human-Vector Interactions in Time and Space

Tzai-Hung Wen [1], Niann-Tai Chang [2], Hwa-Jen Teng [3]

1. Department of Geography, National Taiwan University
2. Department of Plant Medicine, National Pingtung University of Science and Technology
3. Research and Diagnostic Center, Centers of Disease Control, Taiwan

Abstract: In order to depict the population dynamics of dengue vector in detail, a total of 5180 mosquito surveillance spots were set up to collect weekly mosquito data, including number of mosquito eggs and larvae of *Aedes aepypti* and *Aedes albopictus,* in the Kaohsiung City and seven major townships with intensive *Aedes aepypti* activities in the Kaohsiung County from the 7th June, 2009 to 13th June, 2010. We used Google Map as the platform to establish a *GIS-based Mosquito Surveillance Mapping System* for providing health authorities with continuous monitoring of mosquito population dynamics spatiotemporally. Areas with high abundance of mosquitoes in specific periods can be identified as spatial target areas for dengue prevention and control. In addition, spatial-temporal diffusion patterns of dengue cases were also analyzed to capture transmission dynamics of the dengue epidemic in 2009. Analyzing the dynamics of dengue cases and mosquitoes simultaneously can delineate more clear patterns of human-mosquito interactions in time and space and provide more geospatial insights into epidemiological characteristics of dengue transmission.

附錄二（文章手稿）

第一篇

The evaluation of different sampling methods for monitoring *Aedes* vectors in dengue high-risk areas

Abstract

Aedes aegypti and *Ae. albopictus* are the main vectors of dengue in Taiwan. The indicators generated by various sampling methods to estimate the local population of *Aedes* mosquitoes are critical for evaluating the disease-risk and the effectiveness of control measures. Five sampling methods, traditional *Stegomyia* indece (Breteaux index, house index and adult index), modified ovitraps, BG-Sentinel^{TM} mosquito traps and CDC backpack aspirators, were compared for one year in Fensheng City, Kaohsiung County and Chichen District, Kaohsiung City. Monthly fluctuation of all indice (except for BG-Sentinel trap index) presented similar pattern, which were concurred with temperatures and rainfalls of the same year. BG-Sentinel traps collected some number of *Aedes* mosquitoes except for March and April. Breteau index, house index, ovitrap index and backpack aspirator index had significantly higher positive correlation with each other ($r>0.91$, $p<0.000$). BG-Sentinel trap index had significantly positive correlations with *Stegomyia* indice ($r=0.581$-0.6793, $p<0.05$). Additionally, the number of indigenous dengue cases had 2 month time-forward effect with Breteau index, house index, and backpack aspirator index ($r=0.6668$-0.8053, $p<0.05$). In conclusion, mosquito control measures should be conducted 1-3 months before the cases was found and BG-Sentinel^{TM} trap was a sensitive collection method, even in a low population.

Key words: *Aedes*, BG-sentinel trap, CDC Backpack Aspirator, ovitrap

1. Introduction

The *Aedes* mosquitoes are the primary disease vector for dengue, yellow fever, and chikungunya. They are found in tropical and subtropical regions throughout the world. In Taiwan, *A. aegypti* and *A. albopictus* are considered the main dengue transmitter and evidenced the dengue virus were existed in the mosquitoes which were detected by real-time RT-PCR (Yang et al., 2010). The distribution of the former species is geographically limited to southern Taiwan (Chang et al., 2007), and the later species is distributed throughout the island, below an elevation of 1,500m above sea level.

In Taiwan, the dengue case started by import from abroad in early summer, spread out locally, and ended in the winter. The dengue cases and dengue hemorrhagic fever (DHF), which included imported and local cases were 145 to 5,388 and 2 to 242 separately during 2001-2009 in Taiwan and were epidemic annually (CDC database, Taiwan). The type 1 to 3 of dengue viruses were mainly imported to Taiwan and widespread based on different years during 2001-2009 (Lin et al., 2010; CDC database, Taiwan). Specially, the dengue outbreak in 2002 caused 5,336 local cases and 241 DHFs, and the type 2 dengue virus were popular. There were some correlation between *Aedes* mosquito densities and dengue incidence (Scott and Morrison, 2003; Tuan et al., 2009). The diseases which occurred and spread was usually accompanied with increased mosquito densities or thresholds of mosquito density. Therefore, the monitor of mosquito densities based on good sampling methods was the important issue for disease control and spread.

In the present day, the measurement of mosquito densities are mainly based on traditional *Stegomyia* indeces, but the larvae usually hide in the any varied of temporal assorted water-holding containers. The traditional index were easily influenced due to hidden container numbers and considered presenting delayed time to the risk of disease spread. Various kinds of ovitaps including traditional ovitraps with paddles or cotton fabric (Lenhart et al., 2005), and lethal traps (Ritchie et al., 2008; Ritchie et al., 2009; Rapley, et al., 2009; Perich et al., 2003) are appropriate methods for the surveillance of larvae and eggs.

Adult *Aedes* mosquitoes are intimate with humans and activate in the daytime (Harrington et al., 2001). Female *Aedes* mosquitoes are the main vector to dispread the diseases. Its blood feeding is easily interrupted allowing multiple individuals to be bitten in a single gonotrophic cycle (Platt et al., 1997). This behavior also caused the risks of disease spread. There are relationships between the quantities of adult mosquitoes and disease-risk (Scott and Morrison, 2003). The good methods for adult mosquito surveillance are backpack aspirators (Vazquez-Prokopec et al., 2009), sticky ovitraps (Ritchie et al., 2003; Facchinelli et al., 2007; Zhang and Lei, 2008),

and BG-sentinel traps (Maciel-de-Freitas et al., 2006).

The backpack aspirators were considered as most effective way to collect large number mosquitoes (Morrison et al., 2008). Sticky traps are good method to collect blooded adult female mosquito based on oviposition behavior (Ritchie et al., 2003). The BG-sentinel trap utilized the odor to attract and capture female *Aedes* mosquitoes. This effect of method is better than backpack aspirators, sticky ovitraps and other types of traps (Williams et al., 2006; Krockel, et al., 2006; Farajollahi, et al., 2009; Bhalala and Arias, 2009).

In the previous papers, these methods had been compared the efficiency and appropriation, but these methods were not long-term evaluated simultaneously. This study was to evaluate the different methods, backpack aspirators, BG-sentinel trap, ovitrap, and traditional index, for monitoring *Aedes* mosquitoes for one year surveillance in southern Taiwan and their correlations.

2. Materials and Methods

2.1. Study sites

Fengshan City Kaohsiung County (population: 309,062 people; population density, 11,556 people/km^2; households, 94,501;the area, 26.7451 km^2) and Cianjhen District Kaohsiung City (population, 201,477 people: population density, 10,531; household, 61,207; the area, 19.1309 km^2) in Southern Taiwan were chosen as monitored areas (Fig. 1) according to 2007 household registration data. These two sites are high-risk areas of dengue. In these two areas, the dengue case were ever up to 1,741 and 967 cases and dengue hemorrhagic fever were up to 80 and 50 cases separately in 2002.

2.2. Sampling methods used in this study

This surveillance plan executed for one year, from June, 2009 to June, 2010. The sampling methods used in this study were described bellow.

2.2.1. Traditional *Stegomyia* index

The prolonged monitoring by local governments produced the traditional *Stegomyia* index, including Breteau index (the number of positive containers per 100 houses), house index (the percentage of houses infested with larvae and/or pupae), container index (the percentage of water-holding containers infested with active immatures), and adult *Aedes* mosquito index (the average number of *Aedes* adult female per house). These indeces were used to compare the correlation with other indeces of other sampling methods.

2.2.2. Ovitraps

The modified ovitraps (Fig. 1) which were designed similar to HongKong and Hawaii ovitraps were produced by local manufactory, Taiwan. Based on 2 traps (one indoor and one outdoor) for every 400 people, the total number of ovitraps are estimated to be 1,282. Three treatment types in the ovitraps were paddles, or sticky papers, or both all. All the ovitraps contained almost 70-80% volume water which containing growth regulator agents, pyriproxyfen (Chung His Chemical Plant, Ltd) to prevent adult mosquitoes emerging. The number of eggs on paddles (length x width: 15 x 1.5 cm; the middle part of two sides were rugged) and adult mosquitoes on arc-like sticky papers (upper length 27.5cm, lower length, 23.8cm, width, 7 cm) in the ovitraps which were collected weekly were counted and identified the species of mosquitoes.

2.2.3. BG-Sentinel^{TM} mosquito traps and CDC backpack aspirators

Fengshan City Kaohsiung County and Cianjhen District Kaohsiung City were randomly selected four villages for each month. These villages were investigated the mosquito species and quantities by using BG-Sentinel^{TM} mosquito traps (Biogents, Germany) and CDC backpack aspirators (Model 1412, John W. Hock, Gainesville, FL). Sixteen BG-Sentinel^{TM} mosquito traps were positioned eight houses separately for each month (one indoor and one outdoor of each house). The whole BG-sentinel traps were placed the lure (BG lure) and half of traps were combined with carbon dioxide to attract the adult mosquitoes. The trapping time was about 7-8 hours one day (period about from a.m. 8:00 to p.m. 6:00). To the CDC backpack aspirators, there were two groups (two people in one group) to survey 10 to 15 households for each village in the morning, separately.

2.2.4. Statistical analyses

The Pearson correlation was used to test relationships between different methods for *Aedes* populations. The statistic software, STATISTICA 6.1 (StatSoft, Inc, USA) was used to analyze the correlation.

3. Results

3.1. The correlations among different sampling methods

The correlations among different sampling methods were shown as Table 1. The ovitrap index, BG-Sentinel trap index, and CDC Backpack aspirator index mean the percentage of positive households detected by these methods, separately. There are the highest correlations ($r=0.9973$, $p<0.05$) between Breteau index and house index. The house index and Breteau index possess higher positive correlation with

ovitrap index (r=0.9184-0.9246, p<0.05) and CDC backpack aspirator index (r=0.9309-0.9383, p<0.05) and lower positive correlation with BG-Sentinel trap index (r=0.6529-0.6793, p<0.05). The ovitrap index also has high positive correlations with backpack aspirator index (r=0.9785, p<0.05), besides adult *Aedes* mosquito index and local dengue cases. The BG-Sentinel trap index have lower positive correlations with the most other indeces (r=0.5805-0.6793, p<0.05). CDC backpack aspirator index has high positive correlation with most traditional indeces and ovitrap index (r=0.9309-0.9758, p<0.05). Adult *Aedes* mosquito index and local dengue cases has no correlations with other index (r=-0.009-0.5108, p=0.384-0.977), but adult *Aedes* mosquito index have low positive correlation with backpack aspirator index (r=0.5805, p<0.05).

3.2. Monthly tendencies detected by various sampling methods

The monthly tendencies detected by various sampling methods and temperature and rainfall were shown as Fig. 2. In order to present the fluctuation of lower values of some indeces, the original values of the BG trap index and CDC backpack aspirator index were divided by ten. The tendencies of almost all index present similar fluctuation and achieved highest points at Sep, 2009 beside the BG trap index at July, 2009. After August 2009, the tendencies gradually decreased and increased again at March or April 2010. At August, 2009, there were different degrees of drops for most indeces beside Breteau index which still increased.

There were some intervals between local dengue cases and mosquito densities. The shift ahead of local dengue cases were compared the relationships to all the indeces used in this study. The results showed high positive correlations (r=0.5611-0.8053, p=0.001-0.046) among most indeces and 1 to 3 months shift ahead of local dengue cases, besides adult *Aedes* mosquito index and BG-sentinel trap index. The 2 months shift ahead presented higher positive correlations (r=0.6668-0.8053, p=0.001-0.013) than 1 and 3 months shift ahead.

3.3. The species composition and efficiency comparison of various sampling methods

The total numbers and composition of different mosquito species by various sampling methods were showed as Table 2. *Ae. aegypti* are the dominant species of *Aedes* spp. in spit of different sampling methods (60.4% in ovitrap, 95.0% in BG-sentinel trap, and 91.0% in backpack aspirator of total *Aedes* spp. numbers). *Ae. albopictus* are the minus species of *Aedes* spp. (12.1% in ovitrap, 16.1% in BG-sentinel trap, and 10.1% in backpack aspirator of total *Aedes* spp. numbers). The percentage of *Aedes* to total mosquito numbers are 9.5% in ovitrap, 48.2% in

BG-sentinel trap, and 29.4% in backpack aspirator. Other species is *Culex quiquefasciatus* (90.3% in ovitrap, 35.8% in BG-sentinel trap, and 60.4% in backpack aspirator of total mosquito numbers), *Armigeres subalbatus* (0.2% in ovitrap, 16.1% in BG-sentinel trap, and 10.1% in backpack aspirator of total mosquito numbers) and *Cu. fuscanus* (<0.1% in ovitrap).

4. Discussion

There are high positive correlation among most indeces and shift ahead of local dengue cases. Because Breteau index, house index and ovitrap index mainly survey larvae in hided the water-containers, it need 1-2 months for mosquitoes to spread viruses, including mature, stealing blood, spreading virus, and latent period of disease. Therefore, mosquito control measures should be conducted 1-3 months before the cases were found.

The traditional indeces were used continuously for the survey of *Aedes* species in Taiwan. They spend many labor power, time, and money. There are better sampling methods to evaluate the densities of larvae and adult mosquitoes. The ovitraps were designed to let blood-feeding female mosquitoes to lay their eggs due to oviposition behaviors. Our results showed more adult females were captured than males by sticky ovitraps in spit of mosquito species. *Aedes* spp. can be detected and its densities presented similar tendencies with traditional indeces. *Cu. quinquefasciatus* is the dominant species in the ovitrap survey due to its abundance in the southern Taiwan. This result was also observed the paper described Facchinelli et al. 2008. BG-Sentinel traps used the lure to attract *Aedes* spp.. Our results proved higher percentage of *Aedes* spp. were captured (48.2%) and better specificity. This tool is more sensitivity compared to other methods, especially in low densities of *Aedes* spp. (Fig. 2). The backpack aspirator used the attraction power of air to collect any mosquitoes by labor power. *Aedes* spp. was efficiently collected by this method and the proportion of sex was almost the same. Based on similar reason of ovitrap, *Cu. quinquefasciatus* were also mainly captured species.

The BG-Sentinel traps are the most efficient to capture *Aedes* spp. (average 0-10.25 *Aedes* mosquito per household) compared to ovitrap (average 0.0034-0.0836 *Aedes* mosquito per household) and backpack aspirator (average 0-1.95 *Aedes* mosquito per household). This similar result was described by Maciel-de-Freitas et al. 2006. BG-Sentinel traps were restricted to smaller survey regions, household numbers, and costs. But they were more sensitive and specific to detect whatever to *Aedes* species, the quantities or low mosquito densities. There are predominant for ovitraps. They can be disperse large coverage scope, investigate long time

continually and cost low. Although ovitraps detected lowest quantities of average *Aedes* mosquitoes per household, they can detect each month, even low mosquito densities. But the sensitivities were based on large scope survey. Backpack aspirators were mainly based on labor power. The labor power will affect the survey scope. The effect of backpack aspirator lied between that of ovitraps and BG-Sentinel traps.

In conclusion, the effects of various sampling methods used in this study show the high positive correlation to traditional indeces and presented different results for capturing *Aedes* species based on their principles. Most sampling methods showed the high positive correlations to 1-3 month shift ahead of local dengue cases. These methods which compared to traditional indeces showed the same effects for monitoring *Aedes* species and can be evaluated to use continuously for the surveillance of disease-borne vectors.

Acknowledgements

We thank the people who helped collect mosquitoes. This study was supported partly by scientific research grants from the Center for Disease Control, Taiwan in 2009 and 2010 (DOH-98-DC-2037, DOH99-DC-2029).

References

Womack, M. (1993). "The yellow fever mosquito, *Aedes aegypti*". *Wing Beats* **5** (4): 4.

Bhalala, H., Arias, J.R., The Zumba mosquito trap and BG-Sentinel trap: novel surveillance tools for host-seeking mosquitoes. J. Am. Mosq. Control Assoc. 25, 134-139.

Chang, L.H., Hsu, E.L., Teng, H.J., Ho, C.M., 2007. Different survival of Aedes aegypti and Aedes albopictus (Diptera: Culicidae) larvae exposed to low temperatures in Taiwan. J. Med. Entomol. 44, 205-210.

Facchinelli, L., Valerio, L., Pombi, M., Reiter, P. Costantini, C., and Della, T.A., 2007. Development of a novel sticky trap for container-breeding mocquitoes and evaluation of its sampling properties to monitor urban populations of *Aedes albopictus*. Med. Vet. Entomol. 21, 183-195.

Farajollahi, A., Kesavaraju, B., Price, D.C., Williams, G..M., Healy, S.P., Gaugler, R., Nelder, M.P., 2009. Field efficacy of BG-Sentinel and industry-standard traps for *Aedes albopictus* (Diptera: Culicidae) and West Nile virus surveillance. J. Med. Entomol. 46, 919-925.

Harrington, L.C., Edman, J.D., Scott, T.W., 2001. Why do female *Aedes aegypti* (Diptera: Culicidae) feed preferentially and frequently on human blood? J.

Med. Entomol. 38, 411-422.

Krockel, U, Rose, A., Eiras, A.E., Geier, M., 2006. New tools for surveillance of adult yellow fever mosquitoes: comparison of trap catches with human landing rates in an urban environment. J. Am. Mosq. Control Assoc. 22, 229-238.

Lin. C.C., Huang, Y.H., Shu, P.Y., Wu, H.S., Lin, Y.S., Yeh, T.M., Liu, C.C., Lei, H.Y., 2010. Charateristic of dengue disease in Taiwan: 2002-2007. Am. J. Trop. Med. Hyg. 82, 731-739.

Maciel-de-Freitas, R., Eiras, A.E., Lourenco-de-Oliveira, R., 2006. Field evaluation of effectiveness of the BG-Sentinel, a new trap for capturing adult *Aedes aegypti* (Diptera: Culicidae). Mem. Inst. Oswaldo. Cruz. 101, 321-325.

Perich, M.J., Kardec, A. Braga, I.A., Portal, I.F., Burge, R., Zeichner, B.C., Brogdon, W.A., Wirtz, R.A., 2003. Field evaluation of a lethal ovitrap against dengue vectors in Brazil. Med. Vet. Entomol. 17, 205-210.

Rapley, L.P., Johnson, P.H., Williams, C.R., Silcock, R.M., Larkman, M., Long, S.A., Russell, R.C., Ritchie, S.A., 2009. A lethal ovitrap-based mass trapping scheme for dengue control in Australia: II. Impact on populations of the mosquito *Aedes aegypti*. Med. Vet. Entomol. 23, 303-316.

Ritche, S.A., Long, S., Hart, A., Webb, C.E., Russell, R.C., 2003. An adulticidal sticky ovitrap for sampling container-breeding mosquitoes. J. Am. Mosq. Control Assoc. 19, 235-242.

Ritchie, S.A., Long, S.A., McCaffrey, N., Key, C., Lonergan, G., Williams, C.R., 2008. A biodegradable lethal ovitrap for control of container-breeding *Aedes*. J. Am. Mosq. Control Assoc. 24, 47-53.

Ritchie, S.A., Rapley, L.P., Williams, C., Johnson, P.H., Larkman, M., Silcock, R.M., Long, S.A., Russell, R.C., 2009. A lethal ovitrap-based mass trapping scheme for dengue control in Australia: I. Public acceptability and performance of lethal ovitraps. Med. Vet. Entomol. 23, 295-302.

Russell, R.C., Ritchie, S.A., 2004. Surveillance and behavioral investigations of *Aedes aegypti* and *Aedes polynesiensis* in Moorea, French Polynesia, using a sticky ovitrap. J. Am. Mosq. Control. Assoc. 20, 370-375.

Scott, T.W., Morrison, A.C., 2003. *Aedes aegypti* density and the risk of dengue virus transmission. In: Takken W. Scott T.W., editors. Ecological aspects for application of genetically modified mosquitoes. Dordrecht, The Netherlands: FRONTIS. pp. 187-206.

StatSoft, Inc., 2003. STATISTICA (data analysis software system), version 6. www.statsoft.com.

Tuan, Y.C., Hung, M.N., Lin, L.J., Shih, W.Y., Huang, C.C., Chang, C.C., Chen,

M.J., You, C.Y., 2009. Analysis on dengue vector density survey in Kaohsiung and Pingtung areas of southern Taiwan, 2004-2008. Taiwan Epidemiol. Bull. 25, 462-485.

Williams, C.R., Long, S.A., Russell, R.C., Ritchie, S.A., 2006. Field efficacy of the BG-Sentinel compared with CDC Backpack Aspirators and CO2-baited EVS traps for collection of adult Aedes aegypti in Cairns, Queensland, Australia. J. Am. Mosq. Control Assoc. 22, 296-300.

Yang, C.F., Chen, C.F., Su, C.L., Teng, W.J., Lu, L.C., Lin, C., Wang, C.H., Shu, P.Y., Huang, J.H., Wu, H.S., 2010. Screening of mosquitoes using SYBR Green I-based real-time RT-PCR with group-specific primers for detection of Flaviviruses and Alphaviruses in Taiwan. J. Virol. Methods. 168, 147-151.

Zhang, L.Y., Lei, C.L., 2008. Evaluation of sticky ovitraps for the surveillance of *Aedes* (*Stegomysia*) *albopictus* (Skuse) and the screening of oviposition attractants from organic infusions. Ann. Trop. Med. Parasitol. 102, 399-407.



Fig. 1. The position sites of ovitrap and morphology of oivtrap. (A) The black points (the right picture) showed the setting sites of ovitraps in the Fengshan City Kaohsiung County and Cianjhen District Kaohsiung City in Southern Taiwan (the left picture). (B~D) The morphology of the gray canopy (B), black lid (C), and assembly ovitrap (D).

(A)



(B)



Fig. 2. Monthly tendencies detected by various sampling methods used in this study and weather observation were shown as (A) and (B). (A) The original values of the BG trap index, backpack aspirator index and local dengue case number were divided by ten. The left axis showed the percentage of various index, and the right axis showed the values of Breteau index and local dengue cases. (B) The average temperature and rainfall were presented based each month.

Table 1

The correlations of different sampling methods were used in this study for one year surveillance.

	Breteau index	House index	Adult *Aedes* mosquito index	Ovitrap index	BG-Sentinel trap index	CDC Backpack Aspirator index	Local dengue cases	1 month ahead	2 month ahead	3 month ahead	4 month ahead
Breteau index	r=1.000	**r=0.9973 p=0.000**	r=0.4205 p=0.153	**r=0.9246 p=0.000**	**r=0.6529 p=0.016**	**r=0.9383 p=0.000**	r=0.238 p=0.434	**r=0.5718 p=0.041**	**r=0.7106 p=0.006**	**r=0.668 p=0.013**	r=0.4691 p=0.106
House index	-	r=1.000	r=0.4611 p=0.113	**r=0.9184 p=0.000**	**r=0.6793 p=0.011**	**r=0.9309 p=0.000**	r=0.2508 p=0.409	**r=0.5611 p=0.046**	**r=0.6668 p=0.013**	**r=0.6353 p=0.02**	r=0.4859 p=0.092
Adult *Aedes* mosquito index	-	-	r=1.000	r=0.4834 p=0.094	**r=0.5805 p=0.038**	r=0.5108 p=0.074	r=-0.0719 p=0.815	r=-0.0088 p=0.977	r=0.0384 p=0.901	r=0.3759 p=0.206	**r=0.7163 p=0.006**
Ovitrap index	-	-	-	r=1.000	r=0.4978 p=0.084	**r=0.9758 p=0.00**	r=0.2639 p=0.384	**r=0.6609 p=0.014**	**r=0.8053 p=0.001**	**r=0.6854 p=0.01**	r=0.4890 p=0.09
BG-Sentinel trap index	-	-	-	-	r=1.000	r=0.5397 p==0.057	r=-0.009 p=0.977	r=0.0651 p=0.833	r=0.1924 p=0.529	r=0.3766 p=0.205	r=0.4490 p=0.124
CDC Backpack Aspirator index	-	-	-	-	-	r=1.000	r=0.1902 p=0.534	**r=0.6104 p=0.027**	**r=0.7961 p=0.001**	**r=0.7423 p=0.004**	r=0.5262 p=0.065
Local dengue cases	-	-	-	-	-	-	r=1.000	**r=0.6649 p=0.013**	r=0.0838 p=0.785	r=-0.2291 p=0.451	r=-0.293 p=0.331

Table 2. Species composition of adult mosquitoes collected from June, 2009 to June 2010.

Species	Ovitraps		BG-Sentinel traps		Backpack aspirator	
	Females	Males	Females	Males	Females	Males
Aedes aegypti	679(6%)[a]	457(5%)	170(41%)	95(59%)	173(23%)	169(32%)
Aedes albopictus	189(2%)	38(<1%)	10(2%)	4(3%)	26(3%)	8(1%)
Aedes spp.[b]	298(3%)	219(3%)	0(0%)	0(0%)	0(0%)	0(0%)
Total *Aedes* no.	1,166(11%)	714(9%)	180(43%)	99(62%)	199(27%)	177(33%)
Armigeres subalbatus	22(<1%)	18(<1%)	84(20%)	9(5.63%)	87(11.71%)	43(8.02%)
Culex fuscanus	1(<1%)	0	0	0	0	0
Culex quinquefasciatus	10,246(90%)	7,727(91%)	155(37%)	52(33%)	457(62%)	316(59%)
Total	11,435	8,459	419	160	743	536

[a] showed the mosquito numbers and the amount percentage of total collection mosquitoes.

[b] refer the *Aedes* mosquitoes could not be indentified.

第二篇

A population-stratified ovitrap survey with sticky plastics and wooden paddles for *Aedes* adult and egg collections in surveillance of dengue vectors in dengue high-risk areas of Taiwan

Abstracts (250-300 words)

Background: *Aedes aegypti* (L.) and *Aedes albopictus* Skuse are the main vectors of dengue in Taiwan which coexist in southern Taiwan. The fact that high-cost and long-term programs to control dengue vector population very often do not meet the successful goal suggests that preventing and controlling the disease will require new vector surveillance and control technologies based on field studies of population biology and spatial-temporal distribution of the vectors in their natural environment. We present here a first report on the efforts of a large scale sampling of ovitraps (10,380 ovitraps in 5,190 houses) stratified with population (2 traps every 400 inhabitants). The objectives of this study are using a population-stratified ovitrap survey to monitor the dengue vectors in high dengue-risk areas for one year to study spatial-temporal distribution of *Ae. aegypti*.

Methodology/Principal Findings: A population-stratified ovitrap survey based on egg-collection and adult-collection using a modified ovitrap and an insect growth regulator was evaluated for large-scale sampling in high dengue-risk areas, southern Taiwan. From June 2009 to June 2010, 53 data-collection cycles of 53 weeks were carried out. 10,380 ovitraps were set up into 5,190 households with one ovitrap indoors and one outdoors by 400 people per ovitrap. Three treatments (wood sticks, sticky plastics, or both) were assigned by stratified random sampling with infestation of *Ae. aegypti*.

Conclusion/Significance: A population-stratified ovitrap survey with egg and adult collections simultaneously were effective for measuring spatial-temporal

distribution of *Ae. aegypti*. Using winter distribution map can identify key areas for source reduction of the next year.

Key words: Ovitrap, mosquito surveillance, *Aedes aegypti*, *Aedes albopictus*, *Culex quinquefasciatus*.

Introduction

Aedes aegypti (L.) and *Aedes albopictus* Skuse are the main vectors of dengue in Taiwan which coexist in southern Taiwan (Wu et al. 2009). These two mosquito species are also capable of transmitting Yellow fever, Chikungunya, and West Nile fever in other countries. The former species is considered as a peridomestic species, which rest mainly indoors and feed mainly on human. The latter species, *Aedes albopictus*, shares partial larval habitats (temporary small artificial containers) with *Ae. aegypti* but females considered more exophilic and bites humans and other animals mainly outdoors. Both species are day-time biters. Without human vaccines or more effective mosquito control methods, dengue increased extensively in both frequency and territorial range. In Taiwan, since a major DENV-2 outbreak (5,336 human cases) occurred in 2002, a dengue outbreak with multiple dengue serotypes has occurred every year with the scale (86-2,000 human cases) depending on the appropriate control measures or the weather. The control measures include insecticide space spray with law enforcement as well as source reduction.

Dengue vaccine is still under development and will not be available in the near future. The only method to effectively control dengue is to control its mosquito vectors, especially *Ae. aegypti*. However, the most sensitive indicator to reflect recent infection was the number of *Ae. aegypti* females per person (Rodriguez-Figueroa et al. 1995). The diurnal behavior of this species and breeding in small water-holding containers, especially in those are temporary or located inside

the houses, which limit the availability of collection methods for large-scale sampling. Given the biological characters of *Aedes* populations, trapping eggs or adult mosquitoes would be amore appropriate way of locating sites and times where action should be concentrated to prevent or lessen the intensity of disease outbreaks. Hence, baited traps (Morrison et al. 2008), modified traps (Lenhart et al. 2005), lethal ovitraps treated with insecticides (Ritchie et al. 2008, Williams et al. 2007), and sticky traps (Facchinelli et al. 2008, Ritchie et al. 2003) were evaluated their efficacy to collect and control these diurnal container breeders, *Aedes* mosquitoes. Most surveys were conducted either in a limited time or space.

Peridomestic space spraying with insecticide on dengue transmission was reviewed for its efficacy and they concluded this measure is more likely best applied as part of an integrated vector management strategy (Esu et al. 2010). Routine entomological surveillance of dengue program using ovitraps was conducted in Singapore, Hawaii, and Hong Kong based on survey areas. Focks et al. (2003) reviewed all mosquito indexes and concluded that was the most sensitive indicator for predict dengue risk.

The fact that high-cost and long-term programs to control dengue vector population very often do not meet the successful goal suggests that preventing and controlling the disease will require new vector surveillance and control technologies based on field studies of population biology and spatial-temporal distribution of the vectors in their natural environment as well as the susceptible hosts (human population). We present here a first report on the efforts of a large scale sampling of ovitraps (10,380 ovitraps in 5,190 houses) stratified with population (2 traps every 400 inhabitants). The objectives of this study are using a population-stratified ovitrap survey to monitor the dengue vectors in high dengue-risk areas for one year to study spatial-temporal distribution of *Ae. aegypti*.

Materials and Methods

Study site

This study site (22°90'-22 °50' N and 120°01'-120°50' E,) includes Kaohsiung City (571,166 premises with 1,525,642 inhabitants) and partial areas of Kaohsiung County (235,051 premises with 667,369 inhabitants), which are the main high dengue-risk areas with high infestation of *Ae. aegypti* in Taiwan. In general, the site has a tropical climate with an average temperature of 24.7°C (range, 18.8–28.9°C) and separate rainy and dry seasons. The rainy season (averaging 1567.2mmper season) is from May to September, and the dry season (averaging 248.2 mm per season) is from October to April.

Study design and data collection

A population-stratified ovitrap survey with every 400 inhabitants per 2 ovitraps (range, ovitraps per m^2) was conducted. A total of 10,380 ovitraps (one for indoors and outdoors, respectively) were set up into 5,190 households (0.7%) with 2 ovitraps per 400 people (Fig. 1). Three treatments (wooden paddles, sticky plastics, and both) were randomly assigned to each trap of two areas which were stratified by the proportion of *Ae. aegypti* in a total of dengue vectors (*Ae. aegypti* and *Ae. albopictus*). The proportion of *Ae. aegypti* over the total numbers of dengue vectors in the first and second stratified areas were over 90% and 20%-90%, respectively. Ovitraps were set up on the ground in living room of each house and in the shading areas of the outdoors from June, 2009-June 15, 2010.

Ovitrap description

The ovitrap used in Hong Kong (Lee and Fok 2008) and Hawaii () was further modified using materials available locally. The basic design of the ovitrap (Fig. 2) consisted of: 1) a black plastic cup (6.5 cm and 9.0 cm of bottom and top diameter x 13.5 cm high); 2) a polystyrene cover with four 2 cm x 2 cm openings; 3) an gray

umbrella to prevent raining and dusts. The water level of each ovitrap was 60% of the total volume (400 mL). The inside of the cup was either lined with a layer of an inexpensive sticky plastic (24 cm and 27.5 cm of bottom and top x 7 cm high), a thin paddle of wood (a tone depressor, 15 cm long x 1.8 cm wide) or both. The surfaces of the wood paddles were roughed by a machine before use. The wooden paddle as well as sticky plastic was labeled on its posterior with the house number, location of the ovitrap (indoor or outdoor) and treatment type using a permanent marker.

Permission was obtained from inhabitants before the setup of ovitraps. The indoor ovitraps were placed in low (<2 m high) and dark corners of living room while the outdoor ovitraps were setting or hung in low and hidden places outside the houses. The distance between the ovitraps was determined by the availability of viable locations for their placement, which was highly relied on house style and design. The wooden paddles and sticky plastics were collected in individual plastic bags and replaced weekly. Additionally, the water level was checked every week and refilled by tap water, if necessary.

Density map

Statistical analyses

All statistical analyses were performed using Statistica version 5.0 statistical package (StatSoft Holdings, Inc., Tulsa, Oklahoma). The numbers of eggs and adults were analyzed by a randomized completely block design. Statistical analysis was done after log transformation of adult mosquito and egg counts to normalize the distribution and minimize the standard error. Means separation was made using a least significant difference test. Additionally, a paired t test was used to compare the differences in the mean number of *Aedes* mosquitoes collected indoors and outdoors.

Results

The mean number of eggs collected per ovitrap (0.96 eggs per ovitraps, 2.63%)

in wood paddle alone were significantly higher than those (0.08 eggs per ovitrap, 0.5%) of the wood paddle with sticky traps (Table 1). More *Aedes* adults were collected in sticky plastics (2.95 adults per ovitrap, 2.28%) with wood paddles than those collected in sticky plastics alone (1.90 adults per ovitrap, 1.63%). The predominant mosquito species captured included *Ae. aegypti*, *Ae. albopictus*, *Culex quinquefasciatus* Say, and *Armigeres subalbatus* (Coquillett). The weekly fluctuation of these 4 mosquito species were presented as Fig. 3. *Aedes aegypti* oviposited year round with a significantly drop between the 4th week and 19th week. *Culex quinquefasciatus* was a winter species in this area, which had lowest density between the 34th week and the 53th week. Other few mosquitoes collected were *Anopheles sinensis* Wiedemann and *Culex sitiens* Wiedemann. The weekly change of *Ae. aegypti* females collected were highly correlated with larval density of BI (r=0.6438, P=0.000), HI (r=0.6567, *P*=0.000), and Adult index (r=0.6452, *P*=0.000) while the change of *Ae. albopictus* females were not (r=-0.0084-0.2049, *P*>0.05).

One hundred and thirty eight wards were identified by the winter density map for the on-coming source reduction campaign of the next year (Fig. 4 C).

Cost analysis

The budget of this project included financial fee for temporary workers (over) to collect sticky plastics and wood paddles weekly, 10,380 ovitraps production fees (US$1.68), sticky plastics (US$ 0.2727 per sticky plastic).

Discussion

Significantly more *Aedes* adults were collected in sticky plastics with wooden paddles than sticky alone while less eggs were collected in wooden plastic with sticky plastics than wooden paddles alone.

Larger than 4 traps per house were suggested for optimizing ovitrap use for *Ae. aegypti* (Williams et al 2006). However, in our study, only 2 ovitraps per house were

used.

Morrison et al. (2008) proposed a new paradigm for control should include focused surveillance and strategies that kill adult mosquitoes.

If the surveillance responses timely by reducing the trapping size, this will be a sensitive indicator for dengue prevention.

Although the egg yield of ovitraps using wooden paddles has been previously demonstrated to be high (Chadee et al 1995, Lenhart 2005), our results showed a comparatively low egg yield. This may due to not uniform rough surface of wooden paddles. This can be overcome by using a better attractive material (such as sugar cane board). Moreover, our results showed that significantly more *Aedes* adults were collected in sticky plastics with wooden paddles than sticky alone while less eggs were collected in wooden plastic with sticky plastics than wooden paddles alone. This indicated that the wooden paddles provided females to stay inside the ovitraps longer time and were caught by sticky plastics later.

Because of the shortcomings (high cost, insecticide resistance of target vector species, public rejection, and environmental negative impacts) of insecticide spray with large scale, removal trapping , the use of visual, olfactory, and tactile attractants to lure vectors to a small areas where they are killed, become well-attention for alternative vector control or part of integrated vector management (Day and Sjogren 1994). In our study, the highest number of mosquitoes attracted in one trap was 150 *Culex quinquefasciatus* while the highest number of *Ae. aegypti* in one trap was 20.

References

Ai-leen GT, Song RJ (2000) The use of GIS in ovitrap monitoring dengue control in Singapore. Dengue Bull 24: 110-116.

Chadee DD, Corbet PS, Talbot H, et al. (1995) Proportions of eggs laid by *Aedes aegypti* on different substrates within an ovitrap in Trinidad, West Indies. Med Vet Entomol 9: 66-70.

Day JF, Sjogren RD (1994) Vector control by removal trapping. Am. J. Trop. Med. Hyg. 50 suppl.:126-133.

Eisen L, Lozano-Fuentes S (2009) Use of mapping and spatial and space-time modeling approaches in operational control of *Aedes aegypti* and dengue. PLoS Neg Trop Dis 3(4): e41. doi:10.1371/journal.pntd.0000411

Esu E, Lenhart A, Smith L, Horstick O (2010) Effectiveness of peridomestic space spraying with insecticide on dengue transmission; systematic review. Trop Med Inter Health 15: 619-631.

Facchinelli L, Koenraadt CJM, Fanello C, Kijchalao U, Valerio L, Jones JW, Scott TW, and Torre A. Evaluation of a sticky trap for collection *Aedes* (*Stegomyia*) adults in a dengue-endemic area in Thailand. Am. J. Trop. Med. Hyg. 78, 2008: 904-909.

Favier C, Schmit D, Muller-Graf CDM, Cazelles B, Degallier N, et al. (2005) Influence of spatial heterogeneity on an emerging infectious disease: the case of dengue epidemics. Proc Biol Sci 272:1171-1177.

Focks DA. (2003) A review of entomological sampling methods and indicators for dengue vectors, TDR/IDE/Den/03.1. 36 pp.

Lee MW, Fok MY. (2008) Dengue vector surveillance in Hong Kong-2007. Dengue Bull 32: 38-43.

Lenhart AE, Walle M, Cedillo H, Kroger A. (2005) Building a better ovitrap for detecting *Aedes aegypti* oviposition. Acta Tropica 96: 56-59.

Lin CC, Huang YH, Shu PY, Wu HS, Lin YS et al. (2010) Characteristic of Dengue Disease in Taiwan: 2002–2007. Amer J Trop Med Hyg, 82: 731–739.

Morrison AC, Zielinski-Gutierrez E, Scott TW, Rosenberg R (2008) Defining challenges and proposing solutions for control of the virus vector *Aedes aegypti*. PLoS Med 5: e68.

Ritchie SA, Long S, Hart A, Webb CE, Russell RC. (2003) An adulticidal sticky

ovitrap for sampling container-breeding mosquitoes. J Amer Mosq Control Asso 19: 235-242.

Ritchie SA, Long SA, McCaffrey N, Key C, Lonergan G et al. (2008) A blodergradable lethal ovitrap for control of container-breeding *Aedes*. J Amer Mosq Control Asso 24: 47-53.

Ritchie SA, Long SA, Smith G, Pyke A, Knox TB. (2004) Entomological investigations in focus of dengue transmission in Cairns, Queensland, Australia, by using the sticky ovitraps. J Med Entomol 41: 1-4.

Rodriguez-Figueroa L et al. (1995) Risk factors for dengue infection during an outbreak in Yanes, Puerto Rico in 1911. Amer J Trop Med Hygi 52:496-502.

Williams CR, Long SA, Russell RC, Ritchie SA. (2006) Optimizing ovitrap use for *Aedes aegypti* in Cairns, Queensland, Australia: effects of some abiotic factors on field efficacy. J Amer Mosq Control Asso 22: 635-640.

Williams CR, Ritchie SA, Long SA, Dennison N, Russell RC. (2007) Impact of a bifenthrin-treated lethal ovitrap on *Aedes aegypti* oviposition and mortality in North Queensland, Australia. J Med Entomol 44: 256-262.

Wu JW, Teng HJ, Lin C, Wang CY, Liu DP et al. (2009) Recent distribution of vector mosquitoes and epidemiology of the diseases they transmitted in Taiwan. Med Entomol Zool 60:241-252.



Fig. 1 The ovitrap setting in Kaohsiung City and Kaohsiung County.



Fig. 2. The diagram of an ovitrap comprised of 400 ml plastic container in 13.5 height with 9 cm dia. opening and 6 cm Dia. base (A), covered by a black cover (10 cm dia.) with four 1.5 cm dia. openings (B), and a grey-color umbrella-shape lid on top (C) to prevent rain and dust. Inside the ovitrap, a sticky plastic (D), a wood paddle (E) or both was placed.

(A)



(B)



(C)



Fig. 3. The weather (A) and the weekly fluctuation of female mosquitoes in ovitraps (B), Breteaux index, House index, and Adult index (C) by door to door routine surveys conducted in Kaohsiung City and partial Kaohsiung County.



Fig. 4. Dengue case map (A) and density map of *Aedes aegypti* in the whole year (B) and in winter month (C) in Kaoshiung City and partial Kaoshiung County.

Table 1. Fecundity and adults number of *Aedes* collected in ovitraps from June 2009 to June 2010 in Kaoshiung City and partial Kaoshiung County.

Treatment	n	Egg no. per ovitrap Mean±SD	Egg positive rate Mean ± SD	*Aedes* no. per ovitrap Mean± SD	*Aedes* adult postive rate Mean±SD
Wood paddle alone	204	0.69a±0.65	2.63a ±2.24	-----	
Wood paddles and sticky plastics	204	0.08b±0.09	0.50b ±0.48	2.95 a±1.36	2.28a±1.27
Sticky plastic alone	204	-----	-----	1.90 b±2.50	1.63b±1.71

*Means within each column having the same letter are not significantly different at P≥0.05.

附錄三（病媒蚊空間分布圖）

一、埃及斑蚊雌蚊 2009 年 7 月至 2010 年 5 月（11 張）月份空間分布圖。



二、埃及斑蚊雄蚊 2009 年 7 月至 2010 年 5 月（11 張）月份空間分布圖。





三、白線斑蚊雌蚊 2009 年 7 月至 2010 年 5 月（11 張）月份空間分布圖。





五、白線斑蚊雄蚊 2009 年 7 月至 2010 年 5 月（11 張）月份空間分布圖。