


Summary

According to the laboratory surveillance¹, Coxsackie A was the predominant virus type in the past four weeks (Figure 1), and EV71 and EV-D68 virus were active in community. During week 31, the total number of outpatient and ER visits for enterovirus infection was 8,406, which was lower than the threshold (11,000 visits) and showed a downward trend recently, as shown in Figure 2.

Between August 1 and 7, no new EVSC² case was confirmed. There have been eight EVSC cases since 2023 (Figure 3), and cases caused by Coxsackie A4, A6, EV-D68, and Echovirus 21(Figure 3). All cases were below 5 years old (Table 1).

For further information, please visit the Taiwan National Infectious Disease Statistics System (NIDSS) website at https://nidss.cdc.gov.tw/en/Home/Index?op=1





 In terms of the surveillance systems in Taiwan, please see: Jian, S. W., Chen, C. M., Lee, C. Y., & Liu, D. P. (2017). Real-Time Surveillance of Infectious Diseases: Taiwan's Experience. Health security, 15(2), 144-153.
EVSC: Enteroviruses infection with Severe Complications







Figure 2. Trend of outpatient and ER visits for enterovirus infection, 2022-2023



Taiwan CDC 2023

Figure 3. Trend of EVSC, 2022-2023




Table 1. Age and sex distribution of EVSC cases in 2023

Figure 4. Geographical distribution of EVSC cases in 2023