

### Summary

According to the laboratory surveillance<sup>1</sup>, Coxsackie A was the predominant virus type in the past 4 weeks (Figure 1); both numbers of echovirus 11 (ECHO 11) and EV71 cases remained low recently, and all cases were sporadic with mild symptoms. During week 37, a total number of outpatient department and ER visits for enterovirus infection were 7,732. The number of medical visits for enterovirus infection was increasing but below the national baseline (Figure 2).

In week 37, no new enterovirus infection with severe complications (EVSC) case was confirmed. Therefore, the total number remained the same at 28 cases (Figure 3), including 8 deaths due to ECHO 11 (7 cases) and Coxsackie B1 (1 case) infection. ECHO 11 and EV71 were the majority virus types of EVSC cases, and other virus types included Coxsackie A4, A9, A16, B1, B2, B3 and B5. Among EVSC cases, 39.3% were patients less than or equal to one month of age (Table 1). Figure 4 showed the geographical distribution of EVSC cases according to their residential areas.

For further information, please visit the Taiwan National Infectious Disease Statistics System (NIDSS) website at <https://nidss.cdc.gov.tw/en/>




Figure 1. Trend of Enterovirus Isolates, 2017-2018

1. In terms of the surveillance systems in Taiwan, please see: Jian, S. W., Chen, C. M., Lee, C. Y., & Liu, D. P. (2017). Real-Time Surveillance of Infectious Diseases: Taiwan's Experience. *Health security*, 15(2), 144-153.




Figure 2. Trend of outpatient department and ER visits for enterovirus infection, 2017-2018




Figure 3. Trend of EVSC, 2017-2018



Table 1. Age and sex distribution of EVSC cases in 2018

Age(year)	Male	Female	Total
≤ 1 month	7	4	11
1-12 month	1	0	1
1	2	1	3
2	3	1	4
3	2	1	3
4	1	2	3
5	2	0	2
6	0	0	0
7-9	1	0	1
≥ 10	0	0	0
<b>Total</b>	<b>19</b>	<b>9</b>	<b>28</b>




Figure 4. Geographical distribution of EVSC cases in 2018

