

# Taiwan Nosocomial Infections Surveillance System

## I. Preface

The "nosocomial infection" is limited to describing infections that acquired in hospitals, while the "healthcare-associated infection" (HAI) generally refers to infections that patients acquire while receiving treatment for medical or surgical conditions. HAIs may occur in all settings of care, including hospitals, long-term care facilities, homecare facilities, or outpatient departments. In order to respond to continuous evolving in the contents of medical services and the expansion of surveillance range, "healthcare-associated infection" instead of "nosocomial infection" was commonly used internationally as well as in the definition of infection surveillance in the acute care settings that published by the US CDC in 2008. To monitor the occurrence of HAIs effectively, to evaluate the epidemiologic trend of HAIs in Taiwan, and to analyze surveillance data using well-recognized indicators, so that all the information could be made use of collectively to serve as important references for policy making, Taiwan CDC had revised and launched the Taiwan Nosocomial Infections Surveillance System (TNIS System) in 2007. Moreover, strengthening in functions and the utility of the surveillance system is continuously going on. TNIS system not only helps to gather demographic data as well as laboratory results of pathogen identified and antimicrobial susceptibility test for each HAI case, but also provides simple analytical function, so that reporting hospitals can analyze their own data on line as a reference in developing quality improvement initiatives.

## II. Objectives

1. Establish the epidemiological database of HAI in Taiwan
2. Discovery of HAI trends
3. Facilitation of inter- and intra-hospital comparisons that can be used for quality improvement activities
4. Assistance for hospitals in developing the appropriate surveillance mechanism that permits timely recognition of infection control problems.

## III. Reporting methods, data analysis, and feedback

TNIS system adopts voluntary reporting, and each hospital may provide their data either through web-based entry or convey their data electronically through interchange platform. The web-based report mechanism mainly serves for the hospitals which lack HAI surveillance system of their own. Hospital staff enters the HAI data on the TNIS system website directly. The other mechanism, conveying

surveillance data electronically through interchange platform, serves for the hospitals which had built their own HAI surveillance system. However, to enable interoperability between hospital information systems (HIS) and TNIS system, infection control practitioner has to work on vocabularies mapping from local to standard codes and hospital information technology staff has to bridge the connection between the two systems and make the electronic data pack in a standard format according to the working instruction issued by Taiwan CDC. Through this mechanism, surveillance data could be routinely transferred from hospital information systems to the TNIS system automatically. This can save the hospital staff a lot of time because they would not need to repeatedly enter the data to both of hospital surveillance system and TNIS system. At present, more than 470 hospitals enrolled in TNIS system. Hospitals may use TNIS system to manage HAI cases and generate individual hospital reports. Also, Taiwan CDC periodically feedback hospitals with national report as a reference for inter- and intra-hospital comparisons, hope to facilitate hospitals to improve their quality in controlling HAIs and to safeguard the wellbeing of healthcare workers and the general public.

#### **IV. Healthcare-associated infection surveillance data analysis content**

1. Number of medical centers and regional hospitals contributing ICU HAI data in this report
2. Distribution of HAI rates by type of location in the ICUs of medical centers and regional hospitals in 2015.
3. Distribution of device-associated infection rates in the ICUs of medical centers and regional hospitals in 2015.
4. Distribution of major sites of HAI in ICU patients from medical centers and regional hospitals in 2015.
5. Common pathogens of HAI for patients in the ICUs of medical centers in 2015.
6. Common pathogens of HAI for patients in the ICUs of regional hospitals in 2015.
7. Antimicrobial resistance proportions of selected pathogens of HAI in the ICUs of medical centers and regional hospitals in 2015.

#### **V. Surveillance method and main results**

All the analytical results in this report were derived from TNIS system database with data updated to 2016/05/18. In 2015, there were 21 medical centers (205 ICU units) and 83 regional hospitals (270 ICU units) reported both HAI cases and the number of patient-days to TNIS system for at least one calendar month (Table 10). The distributions of HAI rate ((number of HAIs/number of patient-days) ×1000‰) in ICUs of medical centers and regional hospitals are shown in Table 11. There were 5,374 episodes of HAI events occurred during

829,519 patient-days in the ICUs of 21 medical centers; the rate of infections was 6.5‰. However, in the ICUs of the 83 regional hospitals, there were 4,275 episodes of HAI events occurred during 863,479 patient-days; the rate of infections was 5.0‰. The HAI rates of ICUs were higher in medical centers than those in regional hospitals by corresponding types of ICU. The distributions of device-associated infection rate in ICUs ((number of device-associated infections/ number of device-days) ×1000‰) are shown in Figure 2. The pooled mean of catheter-associated urinary tract infection (CAUTI) rates was 3.6‰ in medical centers and 2.6‰ in regional hospitals, and the pooled mean of central line-associated bloodstream infection (CLABSI) rates were 4.3‰ and 3.0‰ respectively, the rate of CAUTI and the rate of CLABSI in ICUs of medical centers are higher than those in regional hospitals; the pooled mean of ventilator-associated pneumonia (VAP) rates in regional hospitals is higher than that in medical centers, which are 1.4‰ and 0.9‰ respectively.

The distribution of site-specific HAIs in ICUs is shown in Table 12, with the bloodstream infections topped the list in medical centers (40.9%), followed by urinary tract (35.7%), and pneumonia (10.3%). In regional hospitals, the urinary tract infections topped the list (37.5%), followed by bloodstream infections (32.9%), and pneumonia (19.5%). The common pathogens for HAIs in ICUs are shown in Table 13 and Table 14. The top three pathogens in the ICUs were *Escherichia coli*, *Candida albicans* and *Klebsiella pneumoniae* in both medical centers and regional hospitals. The proportions of antimicrobial resistance among selected pathogens identified from patients in the ICUs with HAIs are shown in Figure 3. In the ICUs of medical centers, the proportion of *Acinetobacter baumannii* isolates those were resistant to carbapenem (CRAB) is 64.3%, the proportion of *K. pneumoniae* isolates those were resistant to carbapenem (CRKP) is 22.5%, the proportion of *Pseudomonas aeruginosa* isolates those were resistant to carbapenem (CRPA) is 17.4%, the proportion of enterococci isolates those were resistant to vancomycin (VRE) is 37.6%, and the proportion of *Staphylococcus aureus* isolates those were resistant to oxacillin (MRSA) is 68.8%. Meanwhile, the antimicrobial resistance proportions of selected pathogens isolated from patients acquired HAIs in the ICUs of regional hospitals were 77.1%, 23.8%, 18.3%, 32.3% and 71.4% for CRAB, CRKP, CRPA, VRE and MRSA, respectively.

## VI. 2015 Data analysis of HAI in the ICUs of medical centers and regional hospitals

**Table 10** Number of medical centers and regional hospitals contributing ICU HAI data in this report, 2015

Hospital level	1 <sup>st</sup> Quarter		2 <sup>nd</sup> Quarter		3 <sup>rd</sup> Quarter		4 <sup>th</sup> Quarter	
	No. of hospitals	No. of HAIs	No. of hospitals	No. of HAIs	No. of hospitals	No. of HAIs	No. of hospitals	No. of HAIs
Medical center	21	1,349	21	1,285	21	1,430	21	1,321
Regional hospital	83	1,069	83	1,057	82	1,112	83	1,072


Note: Data updated to 2016/05/18

**Table 11** Distribution of healthcare-associated infection rates by type of locations in the ICUs of medical centers and regional hospitals, 2015

Hospital level	Type of locations	No. of units <sup>1</sup>	No. of HAIs	Patient -days	HAI Rate <sup>2</sup> (%)	Percentile		
						25th	50th	75th
Medical center	Medical ICU	54 (53)	1,932	253,297	7.6	5.1	7.5	9.8
	Surgical ICU	63 (62)	2,023	273,299	7.4	5.3	7.2	9.8
	Cardiology ICU	16 (15)	381	63,717	6.0	4.2	5.3	7.4
	Pediatric ICU	43 (42)	519	164,507	3.2	2.1	3.1	4.3
	Medical/surgical	29 (26)	519	74,699	6.9	3.7	6.8	10.4
	Total	205 (198)	5,374	829,519	6.5	3.8	6.0	9.0
Regional hospital	Medical ICU	60 (57)	1,183	255,617	4.6	5.1	4.1	5.2
	Surgical ICU	46 (44)	1,078	173,423	6.2	4.2	5.7	7.2
	Cardiology ICU	13 (12)	149	40,766	3.7	2.4	3.4	3.6
	Pediatric ICU	62 (57)	56	55,687	1.0	0	0	1.5
	Medical/surgical	89 (82)	1,809	337,986	5.4	3.2	4.6	6.3
	Total	270 (252)	4,275	863,479	5.0	2.0	3.8	5.7

Note: 1. Units with patient-days<50 are not included in percentile distribution; the number in parentheses is the number of units meeting minimum requirement for percentile distribution.

2. Healthcare-associated infection rate= (number of HAIs/number of patient-days) ×1000‰. For every unit, monthly data was included for analysis only when the patient days and number of HAI cases were both available.



Note: 1. device-associated infection rate= (number of HAIs/number of device-days) ×1000‰;  
 2. UTI, urinary tract infection; BSI, bloodstream infection;  
 3. Units with device-days<50 are not included in percentile distribution; the number in parentheses is the number of units meeting minimum requirement for percentile distribution.

**Figure 2 Distribution of device-associated infection rates in the ICUs of medical centers and regional hospitals, 2015**

**Table12 Distribution of major types of healthcare-associated infection in the ICU patients from medical centers and regional hospitals, 2015**

Types of infection	Medical center		Regional hospital	
	No.	%	No.	%
Urinary tract	1,924	35.7	1,616	37.5
Bloodstream	2,202	40.9	1,420	32.9
Pneumonia	552	10.3	842	19.5
Surgical site	266	4.9	176	4.1
Other	441	8.2	256	5.9
<b>Total</b>	<b>5,385</b>	<b>100</b>	<b>4,310</b>	<b>100</b>

Note: proportion of specific infection type= (number of specific infection type /number of overall infection) ×100%

**Table 13 Common pathogens of healthcare-associated infections in the ICUs of medical centers, 2015**


Pathogens	Types of Infection											
	Total		Urinary tract		Bloodstream		Pneumonia		Surgical site		Others	
	Rank	No.	Rank	No.	Rank	No.	Rank	No.	Rank	No.	Rank	No.
<i>Escherichia coli</i>	1	591	1	395	12	97	7	28	3	43	8	28
<i>Candida albicans</i>	2	537	2	352	7	137	9	10	10	13	9	25
<i>Klebsiella pneumoniae</i>	3	534	6	138	2	236	2	83	3	43	6	34
<i>Pseudomonas aeruginosa</i>	4	477	6	138	9	123	1	127	1	54	5	35
<i>Enterococcus faecium</i>	5	471	4	228	3	201			7	19	11	23
<i>Acinetobacter baumannii</i>	6	459	9	82	1	253	3	75	9	15	6	34
Yeast-like	7	382	3	293	17	49	10	9	12	6	9	25
Other <i>Candida</i> spp. or NOS	8	340	5	160	4	164	23	1	14	5	14	10
<i>Enterobacter</i> species	9	336	10	53	5	154	5	45	2	46	3	38
<i>E.cloacae</i>		268		43		126		36		38		25
Other <i>Enterobacter</i> spp. or NOS		68		10		28		9		8		13
<i>Staphylococcus aureus</i>	10	288	13	18	6	150	6	42	6	24	2	54
Others		1,820		321		980		133		152		234
Total		6,235		2,178		2,544		553		420		540

Note: 1. isolates of the same species of bacteria, regardless of antimicrobial susceptibility pattern, are counted only once per patient per infection. That is, no duplicate isolates are included;  
2. NOS: not otherwise specified

**Table 14 Common pathogens of healthcare-associated infections in the ICUs of regional hospitals, 2015**

Pathogens	Types of Infection											
	Total		Urinary tract		Bloodstream		Pneumonia		Surgical site		Others	
	Rank	No.	Rank	No.	Rank	No.	Rank	No.	Rank	No.	Rank	No.
<i>Escherichia coli</i>	1	561	1	397	7	91	6	37	4	24	10	12
<i>Candida albicans</i>	2	512	2	321	3	121	8	24	8	18	3	28
<i>Klebsiella pneumoniae</i>	3	511	4	155	2	160	3	143	2	28	5	25
<i>Acinetobacter baumannii</i>	4	504	8	85	1	177	2	177	7	20	1	45
<i>Pseudomonas aeruginosa</i>	5	499	3	158	7	91	1	188	1	35	4	27
<i>Staphylococcus aureus</i>	6	275	13	25	4	117	4	84	3	26	6	23
<i>Enterococcus faecium</i>	7	240	6	110	6	97	27	1	9	10	7	22
Other <i>Candida</i> spp. or NOS	8	216	5	127	9	79	15	4	16	2	14	4
<i>Enterobacter</i> species	9	192	10	43	10	70	5	42	4	24	9	13
<i>E.cloacae</i>		141		33		58		20		19		11
Other <i>Enterobacter</i> spp. or NOS		51		10		12		22		5		2
Coagulase negative staphylococci	10	176	12	26	5	102	20	3	9	10	2	35
Others		1,152		365		489		141		81		76
Total		4,838		1,812		1,594		844		278		310

Note: 1. isolates of the same species of bacteria, regardless of antimicrobial susceptibility pattern, are counted only once per patient per infection. That is, no duplicate isolates are included;  
2. NOS: not otherwise specified



- Note: 1. "Antimicrobial resistant %" indicates the % of Isolates with susceptibility tested to be intermediate or resistant to the antimicrobial specified.
2. CRAB: carbapenem (imipenem or meropenem)-resistant *Acinetobacter baumannii*; CRKP: carbapenem (imipenem, meropenem, or ertapenem)-resistant *Klebsiella pneumoniae*; CRPA: carbapenem (imipenem or meropenem)-resistant *Pseudomonas aeruginosa*; VRE: vancomycin-resistant enterococci (*Enterococcus faecalis*, *Enterococcus faecium*...etc.); MRSA: oxacillin-resistant *Staphylococcus aureus*.

**Figure 3 Antimicrobial resistances of selected pathogens of healthcare-associated infections in the ICUs of medical centers(MC) and regional hospitals(RH), 2015**